

MATHIEU'S SERIES: INEQUALITIES, ASYMPTOTICS AND POSITIVE DEFINITENESS

VIKTOR P. ZASTAVNYI

ABSTRACT. Inequalities, asymptotics and, for some specific cases, asymptotical expansions were obtained for generalized Mathieu's series. A connection between inequalities for Mathieu's series and positive definite and completely monotonic functions.

1. INTRODUCTION

In 1890 Émile Leonard Mathieu [1] conjectured that the inequality

$$(1.1) \quad S(t) := \sum_{k=1}^{\infty} \frac{2k}{(k^2 + t^2)^2} < \frac{1}{t^2}, t > 0$$

is valid. The first proof of (1.1) was published in 1952 in Berg [2]. It was mentioned that, in case $0 < t \leq \sqrt{2}$ inequality (1.1) follows from inequalities Schröder [3, c. 259]

$$S(t) < \frac{2}{(1+t^2)^2} + \frac{4}{(4+t^2)^2} + \frac{1}{4+t^2} = \frac{1}{t^2} - \frac{2(8-t^6)}{t^2(1+t^2)^2(4+t^2)^2}, \quad 0 < t \leq 2,$$

and, in case $t \geq \sqrt{2}$ he applied the summation formula of Euler-Maclaurin.

In 1956 van der Corput and Heflinger [4] got the estimation of remainder term in the summation formula of Euler-Maclaurin for sufficiently wide classes of the functions. The inequality (1.1) follows from this estimation for every $t > \frac{\sqrt{42}}{16}$. If $0 < t < 1$, then the function $\frac{x}{(x^2+t^2)^2}$ is strictly decreasing with respect to $x \in [1, +\infty)$ and, consequently, $S(t) < \frac{2}{(1+t^2)^2} + \int_1^{\infty} \frac{2x dx}{(x^2+t^2)^2} < \frac{1}{t^2}$. They also mentioned the error in Emersleben [5] which can be corrected. Emersleben has resulted two statements from which the inequality (1.1) follows: 1) for every natural $t \in \mathbb{N}$ the inequality $t^2 S(t) \leq 1 + \frac{1}{16t^2}$ holds; 2) the function $t^2 S(t)$ is strictly increasing for $t > 0$. The proof of the second statement contains an error. The monotonicity of the function $t^2 S(t)$ was proved in [4].

In 2007 author obtained the following proof of the inequality (1.1). Let $F(x) = \frac{x}{e^x - 1}$, then $F(0) = 1$ and for every $t \neq 0$ the next well-known equality

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{2k}{(k^2 + t^2)^2} &= \sum_{k=1}^{\infty} \frac{1}{t} \int_0^{\infty} e^{-kx} x \sin tx \, dx = \frac{1}{t} \int_0^{\infty} F(x) \sin tx \, dx = \\ &= \frac{1}{t^2} - \frac{1}{t^2} \int_0^{\infty} (-F'(x)) \cos tx \, dx \end{aligned}$$

valid. Using the result of Polya [6, §6, VII], [7, Theorem 4.3.1]: *if f is convex downwards on $[0, +\infty)$ and $f(+\infty) = 0$, then $\int_0^{\infty} f(x) \cos tx \, dx \geq 0$ for $t \neq 0$* . If, in

2000 *Mathematics Subject Classification*. Primary 26D15, 33E20, 65B15; Secondary 42A82.

Key words and phrases. Mathieu's series, inequalities, asymptotics, positive definite and completely monotonic functions.

addition, f is not a piecewise linear function with equidistant nodes then the last integral is strictly greater than zero (see, for example, Zastavnyi [8]). It is easily can be checked that $F'(+\infty) = F^{(2)}(+\infty) = 0$ and $F^{(3)}(x) = \frac{e^x}{(e^x-1)^4} g(x)$, where $g(x) = e^{2x}(3-x) - 4xe^x - (x+3) = -\sum_{n=5}^{\infty} a_n x^n$, $a_n = \frac{2^{n-1}(n-6)+4n}{n!} > 0$, $n \geq 5$. Then, for every $x > 0$ the inequalities $F^{(3)}(x) < 0$, $F^{(2)}(x) > 0$, $F'(x) < 0$ hold. After application the Polya's result to the function $f(x) = -F'(x)$ the inequality (1.1) follows.

In 1957, Makai [9] proved the inequality (1.1) by the following arguments. It is easy to check that, for every $k \in \mathbb{N}$, $t \neq 0$, the next inequalities

$$\frac{1}{k(k-1)+t^2+1/2} - \frac{1}{k(k+1)+t^2+1/2} < \frac{2k}{(k^2+t^2)^2} < \frac{1}{k(k-1)+t^2} - \frac{1}{k(k+1)+t^2}$$

hold. By the summation, we get

$$(1.2) \quad \frac{1}{t^2+a} < \sum_{k=1}^{\infty} \frac{2k}{(k^2+t^2)^2} < \frac{1}{t^2+b}, t > 0,$$

where $a = \frac{1}{2}$ and $b = 0$. The next natural problem appears: to find maximal b and minimal a so that inequality (1.2) valid. In 1982 Elbert [10] formulated hypothesis that in (1.2) $a = \frac{1}{2\zeta(3)}$ can be taken, where $\zeta(s)$ is Riemann ζ -function.

In 1998, Alzer, Brenner and Ruehr [11] proved that in (1.2) $a = \frac{1}{2\zeta(3)}$ and $b = \frac{1}{6}$ can be taken and these constants are exact. Their proof is an elementary consequence of the inequality

$$(1.3) \quad \sum_{k=1}^{\infty} \frac{k}{(k^2+t^2)^3} < \left(\sum_{k=1}^{\infty} \frac{k}{(k^2+t^2)^2} \right)^2, t \geq 0.$$

The inequality (1.3), as unsolved problem, was published in 1997 [12] and proved by Wilkins [13] in 1998. Note that for the calculation of the exact value of $b = \frac{1}{6}$ the first two terms of the asymptotical expansion

$$\sum_{k=1}^{\infty} \frac{2k}{(k^2+t^2)^2} \sim \sum_{k=0}^{\infty} \frac{(-1)^k B_{2k}}{t^{2k+2}}, t \rightarrow +\infty,$$

was used ([10, 14]). Here, $B_n := B_n(0)$ - the Bernoulli numbers, and $B_n(x)$ - the Bernoulli polynomials.

We have to mention ones more exact result. In 1980, Diananda [15] proved the inequality

$$(1.4) \quad \sum_{k=1}^{\infty} \frac{2k}{(k^2+t^2)^{\mu+1}} < \frac{1}{\mu t^{2\mu}}, \mu > 0, t > 0,$$

which can be derived from

$$\frac{2k\mu}{(k^2+t^2)^{\mu+1}} < \frac{1}{(k(k-1)+t^2)^{\mu}} - \frac{1}{(k(k+1)+t^2)^{\mu}}, \mu > 0, t > 0, k \in \mathbb{N}.$$

References on this topic can be found in [16].

2. MAIN RESULTS

In this paper, we consider the following Mathieu's series

$$(2.1) \quad S(t, u, \gamma, \alpha, \mu) := \sum_{k=1}^{\infty} \frac{2(k+u)^{\gamma}}{((k+u)^{\alpha} + t^{\alpha})^{\mu+1}},$$

$$\gamma \geq 0, \alpha > 0, \delta := \alpha(\mu+1) - \gamma > 1, u > -1, t \geq 0.$$

Let us introduce the next series

$$(2.2) \quad \tilde{S}(t, u, \gamma, \alpha, \mu) := \sum_{k=1}^{\infty} \frac{2(-1)^{k-1}(k+u)^{\gamma}}{((k+u)^{\alpha} + t^{\alpha})^{\mu+1}},$$

$$\gamma \geq 0, \alpha > 0, \delta := \alpha(\mu+1) - \gamma > 0, u > -1, t \geq 0.$$

It is obviously, if $\delta > 1$,

$$(2.3) \quad \tilde{S}(t, u, \gamma, \alpha, \mu) = S(t, u, \gamma, \alpha, \mu) - 2^{1-\delta} S\left(\frac{t}{2}, \frac{u}{2}, \gamma, \alpha, \mu\right).$$

In [17], the problem of obtaining the exact inequalities for $S(t, 0, \frac{\alpha}{2}, \alpha, \mu)$ stated. Note that in [18, Theorem 2], the inequality for $S(t, 0, \frac{\alpha}{2}, \alpha, \mu)$ is not true.

The Theorem 2.4 (see also Corollaries 4.2 and 4.3) contains the proof of the inequalities for the series (2.1) and (2.2) for all admissible parameters. An asymptotics, as $t \rightarrow +\infty$, and an asymptotic representation, for $(\gamma, \alpha) \in \mathbb{Z}_+ \times \mathbb{N}$, as a series in the power of $t^{-\alpha(k+\mu+1)}$, $k \in \mathbb{Z}_+$, obtained as simple consequences.

The Bernoulli, $B_n(x)$, and, Euler, $E_n(x)$, polynomials can be defined by the following generating functions (see, for example, [19, §1.13, §1.14])

$$(2.4) \quad \frac{te^{tx}}{e^t - 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} B_n(x), |t| < 2\pi; \quad \frac{2e^{tx}}{e^t + 1} = \sum_{n=0}^{\infty} \frac{t^n}{n!} E_n(x), |t| < \pi.$$

The connection between the Bernoulli and Euler polynomials follows from the next identity

$$\frac{2e^{tx}}{e^t + 1} = \frac{2e^{tx}}{e^t - 1} - \frac{4e^{tx}}{e^{2t} - 1} = \sum_{n=1}^{\infty} \frac{t^{n-1}}{(n-1)!} \frac{2}{n} \left(B_n(x) - 2^n B_n\left(\frac{x}{2}\right) \right), 0 < |t| < \pi.$$

Consequently, $E_{n-1}(x) = \frac{2}{n} (B_n(x) - 2^n B_n(\frac{x}{2}))$, for every $n \in \mathbb{N}$. The Bernoulli and Euler splines defined by the formulas $b_n(x) = B_n(\{x\})$ and

$$e_n(x) = \frac{2}{n+1} \left(b_{n+1}(x) - 2^{n+1} b_{n+1}\left(\frac{x}{2}\right) \right), n \in \mathbb{Z}_+$$

correspondingly.

Theorem 2.1 and 2.2 give the generalization of Euler-Maclaurin formula.

Theorem 2.1. 1. *If, for some $n \in \mathbb{Z}_+$, the function $F \in C^n[0, +\infty) \cap L[0, +\infty)$, $F^{(n)}$ is a function of bounded variation on $[0, +\infty)$ and $F^{(k)}(+\infty) = 0$ for $0 \leq k \leq n$, then for every $\varepsilon > 0$ and $u \geq 0$ the equality*

$$(2.5) \quad \sum_{k=1}^{\infty} F(\varepsilon k + \varepsilon u) = \frac{1}{\varepsilon} \int_0^{\infty} F(t) dt + \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{(k+1)!} B_{k+1}(-u) F^{(k)}(0) + \frac{(-1)^n \varepsilon^n}{(n+1)!} L_n(\varepsilon, u),$$

where $L_n(\varepsilon, u) = \int_0^{\infty} b_{n+1}\left(\frac{t}{\varepsilon}\right) dF^{(n)}(t + \varepsilon u) + \int_0^1 B_{n+1}(-u + ut) dF^{(n)}(t\varepsilon u),$

$$(2.6) \quad |L_n(\varepsilon, u)| \leq \sup_{0 \leq x \leq 1} |b_{n+1}(x)| V_{\varepsilon u}^\infty(F^{(n)}) + \sup_{-u \leq x \leq 0} |B_{n+1}(x)| V_0^{\varepsilon u}(F^{(n)})$$

valid. If, in addition, $F^{(n)}$ is absolutely continuous on $[0, +\infty)$, then $L_n(\varepsilon, u) = o(1)$ for $\varepsilon \rightarrow +0$, uniformly with respect to $u \in [0, a]$ for every fixed $a > 0$.

2. If, for some $q < 0$, $n \in \mathbb{Z}_+$, the function $F \in C^n[q, +\infty) \cap L[q, +\infty)$, $F^{(n)}$ is a function of bounded variation on $[q, +\infty)$ and $F^{(k)}(+\infty) = 0$, for $0 \leq k \leq n$, then, for every $u < 0$, $\varepsilon \in (0, \frac{a}{u})$, the equality (2.5) valid and

$$(2.7) \quad |L_n(\varepsilon, u)| \leq \sup_{0 \leq x \leq 1} |b_{n+1}(x)| V_{\varepsilon u}^\infty(F^{(n)}) + \sup_{0 \leq x \leq -u} |B_{n+1}(x)| V_{\varepsilon u}^0(F^{(n)}) .$$

If, in addition, $F^{(n)}$ is absolutely continuous on $[q, +\infty)$, then $L_n(\varepsilon, u) = o(1)$, for $\varepsilon \rightarrow +0$, uniformly with respect to $u \in [b, 0]$ for every fixed $b < 0$.

Theorem 2.2. 1. If, for some $n \in \mathbb{Z}_+$, the function $G \in C^n[0, +\infty)$, $G^{(n)}$ is a function of bounded variation on $[0, +\infty)$ and $G^{(k)}(+\infty) = 0$ for $0 \leq k \leq n$, then, for every $\varepsilon > 0$ and $u \geq 0$, the equality

$$(2.8) \quad \sum_{k=1}^{\infty} (-1)^{k-1} G(\varepsilon k + \varepsilon u) = \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{2 k!} E_k(-u) G^{(k)}(0) + \frac{(-1)^n \varepsilon^n}{2 n!} l_n(\varepsilon, u) ,$$

where $l_n(\varepsilon, u) = \int_0^\infty e_n\left(\frac{t}{\varepsilon}\right) dG^{(n)}(t + \varepsilon u) + \int_0^1 E_n(-u + ut) dG^{(n)}(t\varepsilon u) ,$

$$(2.9) \quad |l_n(\varepsilon, u)| \leq \sup_{0 \leq x \leq 2} |e_n(x)| V_{\varepsilon u}^\infty(G^{(n)}) + \sup_{-u \leq x \leq 0} |E_n(x)| V_0^{\varepsilon u}(G^{(n)})$$

is valid. If, in addition, $G^{(n)}$ is absolutely continuous on $[0, +\infty)$, then $l_n(\varepsilon, u) = o(1)$, for $\varepsilon \rightarrow +0$, uniformly on $u \in [0, a]$, for every fixed $a > 0$.

2. If, for some $q < 0$, $n \in \mathbb{Z}_+$, the function $G \in C^n[q, +\infty)$, $G^{(n)}$ is a function of bounded variation on $[q, +\infty)$ and $G^{(k)}(+\infty) = 0$, for $0 \leq k \leq n$, then for every $u < 0$, $\varepsilon \in (0, \frac{a}{u})$, the equality (2.8) is valid and

$$(2.10) \quad |l_n(\varepsilon, u)| \leq \sup_{0 \leq x \leq 2} |e_n(x)| V_{\varepsilon u}^\infty(G^{(n)}) + \sup_{0 \leq x \leq -u} |E_n(x)| V_{\varepsilon u}^0(G^{(n)}) .$$

If, in addition, $G^{(n)}$ is absolutely continuous on $[q, +\infty)$, then $l_n(\varepsilon, u) = o(1)$, for $\varepsilon \rightarrow +0$, uniformly with respect to $u \in [b, 0]$ for every fixed $b < 0$.

Theorem 2.3. 1. If, for some $q \leq 0$, the function $F \in C^\infty[q, +\infty)$ and $F^{(n)} \in L[q, +\infty)$, for every $n \in \mathbb{Z}_+$, then, for every fixed $u \geq 0$, but if $q < 0$, then, for every fixed $u \in \mathbb{R}$, the asymptotic representation

$$(2.11) \quad \sum_{k=1}^{\infty} F(\varepsilon k + \varepsilon u) \sim \frac{1}{\varepsilon} \int_0^\infty F(t) dt + \sum_{k=0}^{\infty} \frac{(-1)^k \varepsilon^k}{(k+1)!} B_{k+1}(-u) F^{(k)}(0) , \quad \varepsilon \rightarrow +0 ,$$

is valid.

2. If, for some $q \leq 0$, the function $G \in C^\infty[q, +\infty)$, $G(+\infty) = 0$ and $G^{(n)} \in L[q, +\infty)$, for every $n \in \mathbb{N}$, then for every fixed $u \geq 0$, but if $q < 0$, then, for every fixed $u \in \mathbb{R}$, the asymptotic representation

$$(2.12) \quad \sum_{k=1}^{\infty} (-1)^{k-1} G(\varepsilon k + \varepsilon u) \sim \sum_{k=0}^{\infty} \frac{(-1)^k \varepsilon^k}{2 k!} E_k(-u) G^{(k)}(0) , \quad \varepsilon \rightarrow +0$$

is valid.

Example 2.1. The function $F(t) = 2t^{2\gamma-1}e^{-t^{2\alpha}}$, for every $\gamma, \alpha \in \mathbb{N}$, satisfy to the conditions of Theorem 2.3 for every $q < 0$. Consequently, F has the representation (2.11). Multiplying (2.11) by ε and setting $\varepsilon^2 = x > 0$, the next asymptotical representations

$$x^\gamma \sum_{k=1}^{\infty} 2(k+u)^{2\gamma-1} e^{-(k+u)^{2\alpha} x^\alpha} \sim \frac{\Gamma(\frac{\gamma}{\alpha})}{\alpha} + \sum_{k=0}^{\infty} \frac{(-1)^{k+1} B_{2(\alpha k + \gamma)}(-u) x^{\alpha k + \gamma}}{(\alpha k + \gamma) k!}, \quad x \rightarrow +0,$$

$$x \sum_{k=1}^{\infty} 2(k+u) e^{-(k+u)^2 x} \sim \sum_{k=0}^{\infty} \frac{(-1)^k B_{2k}(-u) x^k}{k!}, \quad x \rightarrow +0$$

take place, for every fixed $u \in \mathbb{R}$. For $u = 0$, first of them can be found, for example, in [27, Example 11.8].

Theorem 2.4. *Let $\gamma \geq 0$, $\alpha > 0$, $\delta := \alpha(\mu+1) - \gamma > 0$ and $g(x) := x^\gamma(x^\alpha+1)^{-\mu-1}$.*

1. *If $(\gamma, \alpha) \notin \mathbb{Z}_+ \times \mathbb{N}$, then $g \in C^r[0, +\infty)$, $g \notin C^{r+1}[0, +\infty)$ and, for every integer $n \in [0, r]$, the function $g^{(n)}$ is absolutely continuous on $[0, +\infty)$, where*

$$(2.13) \quad r = \begin{cases} [\gamma] & , \quad \gamma \notin \mathbb{Z}_+, \\ \gamma + [\alpha] & , \quad \gamma \in \mathbb{Z}_+, \alpha \notin \mathbb{Z}_+. \end{cases}$$

If $\gamma \notin \mathbb{Z}_+$, then $g^{(p)}(0) = 0$ for every integer $p \in [0, r]$. If $\gamma \in \mathbb{Z}_+$, $\alpha \notin \mathbb{N}$, then $g^{(p)}(0) = 0$ for every integer $p \in [0, r]$, $p \neq \gamma$ and $g^{(\gamma)}(0) = \gamma!$.

Furthermore, the function $G = g$ satisfies to the conditions of the statement 1 of Theorem 2.2 for every integer $n \in [0, r]$.

If, in addition, $\delta > 1$, then the function $F = g$ satisfies to the conditions of the statement 1 of Theorem 2.1 for every integer $n \in [0, r]$.

2. *If $(\gamma, \alpha) \in \mathbb{Z}_+ \times \mathbb{N}$, then $g \in C^\infty(-1, +\infty)$ and the function $G = g$ satisfies to the conditions of Theorem 2.2 for every $q \in (-1, 0)$, $n \in \mathbb{Z}_+$, and the function $g^{(n)}$ is absolutely continuous on $[q, +\infty)$. Furthermore, for every fixed $u \in \mathbb{R}$ the next asymptotic representation*

$$(2.14) \quad \tilde{S}(t, u, \gamma, \alpha, \mu) \sim \sum_{k=0}^{\infty} \frac{(-1)^{k(\alpha+1)+\gamma}}{t^{\alpha(k+\mu+1)}} \cdot \frac{\Gamma(\mu+k+1) E_{k\alpha+\gamma}(-u)}{\Gamma(\mu+1)\Gamma(k+1)}, \quad t \rightarrow +\infty$$

is valid. If, in addition, $\delta > 1$, then the function $F = g$ satisfies to the conditions of Theorem 2.1 for every $q \in (-1, 0)$, $n \in \mathbb{Z}_+$. In this case, for every fixed $u \in \mathbb{R}$, the next asymptotic representation

$$(2.15) \quad S(t, u, \gamma, \alpha, \mu) \sim \frac{1}{t^{\alpha(\mu+1)-\gamma-1}} \cdot \frac{2}{\alpha} B\left(\frac{\gamma+1}{\alpha}, \mu+1 - \frac{\gamma+1}{\alpha}\right) +$$

$$\sum_{k=0}^{\infty} \frac{(-1)^{k(\alpha+1)+\gamma}}{t^{\alpha(k+\mu+1)}} \cdot \frac{2\Gamma(\mu+k+1)}{\Gamma(\mu+1)\Gamma(k+1)} \cdot \frac{B_{k\alpha+\gamma+1}(-u)}{k\alpha+\gamma+1}, \quad t \rightarrow +\infty,$$

is valid.

The asymptotic representation (2.14), for $u = 0$, $\gamma = 1$, $\alpha = 2$, and, $\mu = 1$, obtained in [20] by using a different method.

Further, let us consider some generalization of the inequalities (1.2) and (1.4). The next Theorem 2.5 is a sufficient conditions for the existence of double inequalities.

Theorem 2.5. *Let the next conditions be satisfied:*

- i) $S(t) \in C[0, +\infty)$, and $S(t) > 0$, $t \geq 0$.
- ii) *There exist the positive constants $C, \delta_1, A, \beta_1 > 0$ such that*

$$(2.16) \quad S(t) = \frac{C}{t^{\delta_1}} - \frac{A}{t^{\delta_1 + \beta_1}} + o\left(\frac{1}{t^{\delta_1 + \beta_1}}\right), \quad t \rightarrow +\infty.$$

- iii) *For every $t > 0$, $S(t) < \frac{C}{t^{\delta_1}}$.*

Then

1. *If $0 < \beta < \beta_1$, the inequality $S(t) \leq C(t^\beta + b)^{-\frac{\delta_1}{\beta}}$, $t > 0$, is impossible for every $b > 0$.*
2. *If $\beta > \beta_1$, the inequality $S(t) \geq C(t^\beta + a)^{-\frac{\delta_1}{\beta}}$, $t > 0$, is impossible for every $a \geq 0$.*
3. *Let $m = \inf_{t \geq 0} f(t)$, $M = \sup_{t \geq 0} f(t)$, here $f(t) = \left(\frac{C}{S(t)}\right)^{\frac{\beta_1}{\delta_1}} - t^{\beta_1}$. Then $f(+\infty) = \frac{\beta_1 A}{\delta_1 C}$, $0 < m \leq M < +\infty$, and inequality*

$$(2.17) \quad \frac{C}{(t^{\beta_1} + a)^{\frac{\delta_1}{\beta_1}}} \leq S(t) \leq \frac{C}{(t^{\beta_1} + b)^{\frac{\delta_1}{\beta_1}}}, \quad t > 0$$

is valid iff $0 \leq b \leq m$ and $a \geq M$.

Example 2.2. For $S(t) = S(t, u, \gamma, \alpha, \mu)$, $\gamma = 0$, $\alpha > 0$, $\alpha(\mu + 1) > 1$, $u \geq 0$ the conditions of Theorem 2.5 be fulfilled. In this case (see Corollary 4.2) $S(t, u, 0, \alpha, \mu) = \frac{C}{t^{\delta_1}} - \frac{A}{t^{\delta_1 + \beta_1}} + o\left(\frac{1}{t^{\delta_1 + \beta_1}}\right)$ as $t \rightarrow +\infty$, where $C = \frac{2}{\alpha} B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)$, $\delta_1 = \alpha(\mu + 1) - 1$, $A = 1 + 2u$, $\beta_1 = 1$. Furthermore, for every $t > 0$, the inequality

$$S(t, u, 0, \alpha, \mu) < 2 \int_0^\infty \frac{dx}{((x + u)^\alpha + t^\alpha)^{\mu+1}} \leq 2 \int_0^\infty \frac{dx}{(x^\alpha + t^\alpha)^{\mu+1}} = \frac{C}{t^{\delta_1}}$$

is valid. Then, for $m = m(u, \alpha, \mu) = \inf_{t \geq 0} f(t)$, and $M = M(u, \alpha, \mu) = \sup_{t \geq 0} f(t)$, where

$$f(t) = \left(\frac{\frac{2}{\alpha} B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)}{S(t, u, 0, \alpha, \mu)} \right)^{\frac{1}{\alpha(\mu+1)-1}} - t,$$

the next inequalities

$$\frac{\frac{2}{\alpha} B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)}{(t + M)^{\alpha(\mu+1)-1}} \leq \sum_{k=1}^\infty \frac{2}{((k + u)^\alpha + t^\alpha)^{\mu+1}} \leq \frac{\frac{2}{\alpha} B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)}{(t + m)^{\alpha(\mu+1)-1}}, \quad t > 0,$$

$$0 < m \leq \min\{f(0), f(+\infty)\} \leq \max\{f(0), f(+\infty)\} \leq M < +\infty,$$

is valid. Here

$$f(0) = \left(\frac{\frac{2}{\alpha} B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)}{S(0, u, 0, \alpha, \mu)} \right)^{\frac{1}{\alpha(\mu+1)-1}}, \quad f(+\infty) = \frac{\frac{1}{2} + u}{(\mu + 1 - \frac{1}{\alpha}) B\left(\frac{1}{\alpha}, \mu + 1 - \frac{1}{\alpha}\right)}.$$

In the first inequality m and M are the best constants. It is easy to see that $m < M$.

Theorem 2.6. *Let g is convex downwards on $(0, +\infty)$ and $\int_1^{+\infty} g(x) dx$ converges. Then, on the open set $E := \{(u, y) \in \mathbb{R}^2 : (k + u)^2 + y > 0, k \in \mathbb{N}\}$, the function*

$$(2.18) \quad S(u, y) := \sum_{k=1}^\infty 2(k + u)g((k + u)^2 + y), \quad (u, y) \in E,$$

is well define and have the next properties:

1. $S(u, y) \in C(E)$. If $(u_0, y_0) \in E$, then $(u, y) \in E$ for every $y > y_0$ and $S(u_0, +\infty) = 0$. For every fixed $u \geq -1$, the function $S(u, y)$ is nonnegative, convex downwards with respect to $y \in (-(1+u)^2, +\infty)$, and decreases. If, in addition, $g(x) > 0$ for every $x > 0$, then, for every fixed $u \geq -1$, the function $S(u, y)$ is strictly decreasing with respect to $y \in (-(1+u)^2, +\infty)$.
2. For every $u \geq -1$, $u^2 + u + y > 0$, the inequality

$$(2.19) \quad F((1+u)^2 + y) + \left(\frac{1}{2} + u\right) g((1+u)^2 + y) \leq S(u, y) \leq F(u^2 + u + y)$$

is valid, where

$$(2.20) \quad F(t) := \int_t^{+\infty} g(x) dx, \quad t > 0, \quad \left(t \geq 0, \text{ if } \int_0^{+\infty} g(x) dx < +\infty\right).$$

Moreover, the right hand side of (2.19) going to be the equality iff the function g is linear on each segment $[a_k, a_{k+1}]$, $k \in \mathbb{N}$, where $a_k = (k - \frac{1}{2} + u)^2 + y - \frac{1}{4}$. The left hand side of (2.19) valid for every $u \geq -\frac{3}{2}$, $y > -(1+u)^2$ and going to be the equality only if the function g is linear on each segment $[s_k, s_{k+1}]$, $k \in \mathbb{N}$, where $s_k = (k+u)^2 + y$.

Remark 2.1. Let the function g is convex downwards on $(0, +\infty)$, $\int_1^{+\infty} g(x) dx$ converges, and $g(x) > 0$ for every $x > 0$. It follows from (2.19) that, for $u \geq 0$,

$$(2.21) \quad F(a+y) \leq S(u, y) \leq F(b+y), \quad y > 0$$

is valid for $a = (1+u)^2$ and $b = u^2 + u$. In this case, the left hand side of (2.21) is strict inequality. If the inequality (2.21) valid for some $a = a_0 \geq 0$ and $b = b_0 \geq 0$, then it valid for every $a \geq a_0$ and $0 \leq b \leq b_0$, moreover, $b_0 \leq a_0$. Therefore, the problem of definition of the quantities

$$(2.22) \quad \begin{aligned} m(u) &:= \sup\{b \geq 0 : S(u, y) \leq F(b+y), \quad y > 0\}, \quad u \geq 0, \\ M(u) &:= \inf\{a \geq 0 : F(a+y) \leq S(u, y), \quad y > 0\}, \quad u \geq 0 \end{aligned}$$

naturally appears. It is obviously

$$(2.23) \quad u^2 + u \leq m(u) \leq M(u) \leq (1+u)^2, \quad u \geq 0.$$

Because the function F strictly decrease, the inverse function F^{-1} exist such that it is continuous, positive, and strictly decreasing over I , where $I = (0, +\infty)$ if $F(+0) = +\infty$ and $I = (0, F(+0)]$ if $F(+0) < +\infty$. It follows from (2.19) that

$$(2.24) \quad \psi(u, y) := F^{-1}(S(u, y)) - y, \quad (u, y) \in \mathbb{R}_+^2 := [0, +\infty) \times [0, +\infty)$$

is well defined and, consequently, is continuous on $\mathbb{R}_+^2$. Moreover, the next inequalities

$$(2.25) \quad u^2 + u \leq \psi(u, y) < (1+u)^2, \quad (u, y) \in \mathbb{R}_+^2$$

is valid. It is obviously, for fixed $a, b, u \in [0, +\infty)$, the inequality (2.21) is equivalent to the inequality $b \leq \psi(u, y) \leq a$, $y > 0$. Therefore,

$$(2.26) \quad \begin{aligned} m(u) &= \inf_{y \geq 0} \psi(u, y), \quad u \geq 0, \\ M(u) &= \sup_{y \geq 0} \psi(u, y), \quad u \geq 0. \end{aligned}$$

Remark 2.2. Let g is convex downwards on $(0, +\infty)$, $\int_1^{+\infty} g(x) dx$ converges, and $g(x) > 0$ for every $x > 0$. Then:

1) $m(u) = u^2 + u$ for some $u \geq 0 \iff \psi(u, +\infty) = u^2 + u$, or some $y \geq 0$ the functions g is linear on each segment $[a_k, a_{k+1}]$, $k \in \mathbb{N}$, where $a_k = (k - \frac{1}{2} + u)^2 + y - \frac{1}{4}$.

2) $M(u) = (1 + u)^2$ for some $u \geq 0 \iff \psi(u, +\infty) = (1 + u)^2$.

3) If, for some $u \geq 0$, the equality $m(u) = M(u) = c \geq 0$ is valid, then the inequality (2.21) is valid for $a = b = c$. Therefore,
(2.27)

$$m(u) = M(u) = c \geq 0 \iff \sum_{k=1}^{\infty} 2(k+u)g((k+u)^2 + y) \equiv \int_{c+y}^{+\infty} g(x) dx, \quad y > 0.$$

For example, if $g(x) = e^{-\lambda x}$, $\lambda > 0$, then $S(u, 0) = \sum_{k=1}^{\infty} 2(k+u)e^{-\lambda(k+u)^2}$,

$$S(u, y) = e^{-\lambda y} S(u, 0), \quad F(t) = \frac{e^{-\lambda t}}{\lambda}, \quad F^{-1}(s) = -\frac{1}{\lambda} \ln(\lambda s), \quad \psi(u, y) \equiv -\frac{1}{\lambda} \ln(\lambda S(u, 0))$$

and, consequently, $m(u) = M(u) = -\frac{1}{\lambda} \ln(\lambda S(u, 0))$. In this case the inequalities (2.25) is strict. Therefore,

$$(2.28) \quad u^2 + u < -\frac{1}{\lambda} \ln \left(\lambda \sum_{k=1}^{\infty} 2(k+u)e^{-\lambda(k+u)^2} \right) < (u+1)^2, \quad u \geq 0, \quad \lambda > 0.$$

Example 2.3. If $g(x) = x^{-\mu-1}$, $\mu > 0$, $x > 0$, then $F(t) = \frac{1}{\mu t^\mu}$, $F^{-1}(s) = \left(\frac{1}{\mu s}\right)^{\frac{1}{\mu}}$, $t, s > 0$ and, for every $u \geq -1$, $u^2 + u + y > 0$, the inequality (see (2.19))
(2.29)

$$\frac{1}{\mu((1+u)^2 + y)^\mu} + \frac{\frac{1}{2} + u}{((1+u)^2 + y)^{\mu+1}} < \sum_{k=1}^{\infty} \frac{2(k+u)}{((k+u)^2 + y)^{\mu+1}} < \frac{1}{\mu(u^2 + u + y)^\mu}$$

is valid. Left hand side of (2.29) valid for every $u \geq -\frac{3}{2}$, $y > -(1+u)^2$.

In the case $\gamma = 1$, $\alpha = 2$, let us denote the series (2.1) by

$$(2.30) \quad S_\mu(t, u) := S(t, u, 1, 2, \mu) = \sum_{k=1}^{\infty} \frac{2(k+u)}{((k+u)^2 + t^2)^{\mu+1}}, \quad \mu > 0, \quad u, t \in \mathbb{R}.$$

If $-u \in \mathbb{N}$, then, for $k = -u$, the term be absent. Therefore, the function $S_\mu(t, u)$ is continuous in $t \in \mathbb{R}$ for every fixed $u \in \mathbb{R}$. It follows from (2.15) that, for every fixed $u \in \mathbb{R}$, the relations

(2.31)

$$S_\mu(t, u) \sim \sum_{k=0}^{\infty} \frac{(-1)^k B_{2k}(-u)}{t^{2(k+\mu)}} \cdot \frac{\Gamma(\mu+k)}{\Gamma(\mu+1)\Gamma(k+1)}, \quad t \rightarrow +\infty,$$

$$S_\mu(t, u) = \frac{1}{\mu t^{2\mu}} - \frac{u^2 + u + \frac{1}{6}}{t^{2\mu+2}} + \frac{(u^2 + u)^2 - \frac{1}{30}}{t^{2\mu+4}} \cdot \frac{\mu+1}{2} + O\left(\frac{1}{t^{2\mu+6}}\right), \quad t \rightarrow +\infty$$

hold.

Theorem 2.7. For every $t > 0$, $p > 0$, $u > -1$, $\mu > 0$ the inequality

$$(2.32) \quad \left| \frac{1}{\mu(p^2 + t^2)^\mu} - S_\mu(t, u) \right| < \begin{cases} \frac{1}{\mu p^{2\mu}} - S_\mu(0, u) & , \quad p - u \leq \frac{1}{2}, \\ S_\mu(0, u) - \frac{1}{\mu p^{2\mu}} & , \quad p - u \geq 1, \end{cases}$$

is valid.

Theorem 2.8. For $\mu > 0$, $u \geq 0$, let us set $m_\mu(u) = \inf_{t \geq 0} f_{\mu,u}(t)$, $M_\mu(u) = \sup_{t \geq 0} f_{\mu,u}(t)$, where $f_{\mu,u}(t) = (\mu S_\mu(t, u))^{-\frac{1}{\mu}} - t^2$, $t \in \mathbb{R}$. Then $f_{\mu,u}(+\infty) = u^2 + u + \frac{1}{6}$, $0 < m_\mu(u) \leq M_\mu(u) < +\infty$ and inequality

$$(2.33) \quad \frac{1}{\mu(t^2 + a)^\mu} \leq S_\mu(t, u) \leq \frac{1}{\mu(t^2 + b)^\mu}, \quad t > 0$$

is valid iff $0 \leq b \leq m_\mu(u)$ and $a \geq M_\mu(u)$. Moreover, the next statements hold:

1. $u^2 + u < m_\mu(u) \leq u^2 + u + \frac{1}{6} < u^2 + u + \frac{1}{4} < M_\mu(u) < (1 + u)^2$, $\mu > 0$, $u \geq 0$.
2. If $\nu > \mu > 0$, $u \geq 0$, then $m_\nu(u) \leq m_\mu(u)$, $M_\nu(u) \geq M_\mu(u)$.
3. For every $\nu > 0$, $u \geq 0$, there exist $t_{\nu,u} \geq 0$ such that $f'_{\nu,u}(t) < 0$ for every $t > t_{\nu,u}$ and, consequently, the strict inequality

$$(2.34) \quad \frac{1}{\nu(t^2 + f_{\nu,u}(t_{\nu,u}))^\nu} < S_\nu(t, u) < \frac{1}{\nu(t^2 + f_{\nu,u}(+\infty))^\nu}, \quad t > t_{\nu,u}$$

is valid.

4. Let us suppose that, for some $\nu > 0$, $u \geq 0$, the inequality

$$(2.35) \quad ((\nu + 1)S_{\nu+1}(t, u))^{\frac{1}{\nu+1}} \leq (\nu S_\nu(t, u))^{\frac{1}{\nu}}, \quad t \geq 0$$

holds. Then:

- 1) the function $f_{\nu,u}(t)$ strictly decreases in $t \in [0, +\infty)$, the inequality (2.34) valid for $t_{\nu,u} = 0$, $M_\nu(u) = f_{\nu,u}(0)$ and, for every $\mu \in (0, \nu + 1]$, the equality $m_\mu(u) = u^2 + u + \frac{1}{6}$ and strict inequality $S_\mu(t, u) < \frac{1}{\mu(t^2 + m_\mu(u))^\mu}$, $t \geq 0$ take place.
- 2) For every $0 < \mu \leq \nu$, $0 \leq a \leq u^2 + u + \frac{1}{6}$, the inequality $\frac{d}{dt}((t^2 + a)^\mu S_\mu(t, u)) > 0$, $t > 0$ is valid.
5. 1) For every $\mu \in (0, 2]$, the equality $m_\mu(0) = \frac{1}{6}$ and strict inequality $S_\mu(t, 0) < \frac{1}{\mu(t^2 + \frac{1}{6})^\mu}$, $t \geq 0$, take place.
- 2) For every $0 < \mu \leq 1$, $0 \leq a \leq \frac{1}{6}$, $t > 0$, the inequality $\frac{d}{dt}((t^2 + a)^\mu S_\mu(t, 0)) > 0$ is valid.

As we mention in introduction, the equality $m_\mu(0) = \frac{1}{6}$, for $\mu = 1$, proved in [11], and the inequality $\frac{d}{dt}((t^2 + a)^\mu S_\mu(t, 0)) > 0$, $t > 0$ for $\mu = 1$, $a = 0$ proved in [4].

It follows from Theorem 2.8 that, for $u \geq 0$ the relations

$$(2.36) \quad \begin{aligned} m_\infty(u) &:= \inf_{\mu > 0} m_\mu(u) = \lim_{\mu \rightarrow +\infty} m_\mu(u), \\ M_\infty(u) &:= \sup_{\mu > 0} M_\mu(u) = \lim_{\mu \rightarrow +\infty} M_\mu(u), \\ u^2 + u &\leq m_\infty(u) \leq u^2 + u + \frac{1}{6} < u^2 + u + \frac{1}{4} < M_\infty(u) \leq (1 + u)^2 \end{aligned}$$

take place.

The function f is said to be completely monotonic on $(0, +\infty)$ ($f \in M(0, +\infty)$), if $f \in C^\infty(0, +\infty)$ and for every $k \in \mathbb{Z}_+$, $t > 0$, the inequality $(-1)^k f^{(k)}(t) \geq 0$ holds. The Hausdorff-Bernstein-Widder's Theorem state that $f \in M(0, +\infty)$ iff $f(t) = \int_0^{+\infty} e^{-tx} d\mu(x)$, $t > 0$, where μ is nonnegative Borel measure on $[0, +\infty)$ such that the integral converge for every $t > 0$. Besides, the measure μ is finite on $[0, +\infty)$ iff $f(+0) < +\infty$.

Theorem 2.9. *Let the quantities $m_\infty(u)$, $M_\infty(u)$ are defined by (2.36), for $u \geq 0$, and*

$$(2.37) \quad \begin{aligned} \psi_{p,u,\mu}(t) &:= \frac{1}{\mu(p+t)^\mu} - S_\mu(\sqrt{t}, u), \quad p \geq 0, \mu > 0, t > 0, \\ \varphi_u(x) &:= x \sum_{k=1}^{\infty} 2(k+u)e^{-(k+u)^2 x}, \quad x > 0. \end{aligned}$$

Then, for each fixed $u \geq 0$, the next statements hold:

1. $\psi_{p,u,\mu} \in M(0, +\infty)$ for every $\mu > 0 \iff \psi_{p,u,\mu} \in M(0, +\infty)$ for some $\mu > 0 \iff 0 \leq p \leq m_\infty(u)$.
2. $-\psi_{p,u,\mu} \in M(0, +\infty)$ for every $\mu > 0 \iff -\psi_{p,u,\mu} \in M(0, +\infty)$ for some $\mu > 0 \iff p \geq M_\infty(u)$.
3. $m_\infty(u) = -\sup_{x>0} \frac{\ln \varphi_u(x)}{x}$, $M_\infty(u) = -\inf_{x>0} \frac{\ln \varphi_u(x)}{x}$.
4. $u^2 + u < m_\infty(u) \leq u^2 + u + \frac{1}{6}$, $M_\infty(u) = (1+u)^2$.

The next corollary immediately follows from Theorem 2.9.

Corollary 2.10. *If $u \geq 0$ and $0 \leq p \leq u^2 + u$, then $\psi_{p,u,\mu} \in M(0, +\infty)$ for every $\mu > 0$.*

The function $f : \mathbb{R}^m \rightarrow \mathbb{C}$ is said to be positive definite on $\mathbb{R}^m$, $m \in \mathbb{N}$, if for every $n \in \mathbb{N}$, $\{x_k\}_{k=1}^n \subset \mathbb{R}^m$ and $\{c_k\}_{k=1}^n \subset \mathbb{C}$ the next inequality

$$\sum_{k,j=1}^n c_k \bar{c}_j f(x_k - x_j) \geq 0$$

hold. About the positive definite functions see, for example, [21] and references therein. For such a function the next inequality $|f(x)| \leq f(0)$, $x \in \mathbb{R}^m$, holds, and from continuity at zero follows continuity in $\mathbb{R}^m$. By Bochner's Theorem the function f is positive definite and continuous in $\mathbb{R}^m$ iff $f(x) = \int_{\mathbb{R}^m} e^{-i\langle u, x \rangle} d\mu(u)$, where μ is nonnegative, finite, Borel's measure on $\mathbb{R}^m$. From this Theorem immediately follows the criteria of positive definiteness in Fourier's transform term: if $f \in C(\mathbb{R}^m) \cap L(\mathbb{R}^m)$, then f is positive definite on $\mathbb{R}^m$ iff

$$\hat{f}(x) = F_m(f)(x) := \int_{\mathbb{R}^m} e^{i\langle x, y \rangle} f(y) dy \geq 0, \quad x \in \mathbb{R}^m,$$

where $\langle x, y \rangle = x_1 y_1 + x_2 y_2 + \dots + x_m y_m$ - scalar product in $\mathbb{R}^m$, and, in this case $F_m(f) \in L(\mathbb{R}^m)$.

Let us denote by $\Phi(l_2^m)$ a class of continuous functions $f : [0, +\infty) \rightarrow \mathbb{R}$ such that function $f(\|x\|_2)$ is positive definite on $\mathbb{R}^m$. Here $\|x\|_2 = \sqrt{\langle x, x \rangle}$ the Euclidean norm in $\mathbb{R}^m$. Obviously, $\Phi(l_2^{m+1}) \subset \Phi(l_2^m)$ and, by well-known Schoenberg's theorem

$$(2.38) \quad \Phi(l_2) = \bigcap_{m=1}^{\infty} \Phi(l_2^m) = \left\{ f(t) = \int_0^{+\infty} \exp(-t^2 s) d\mu(s), \mu \in P_+ \right\},$$

where P_+ - the set of all finite, nonnegative, Borel measures on $[0, +\infty)$. It follows from the Schoenberg's and Hausdorff-Bernstein-Widder's theorems that $f \in \Phi(l_2)$ iff $f \in C[0, +\infty)$ and $f(\sqrt{x}) \in M(0, +\infty)$.

Theorem 2.11. *Let $m_\infty(u)$, $M_\infty(u)$ are defined, for $u \geq 0$, by (2.36) and $g_{p,u}(x) := \frac{e^{-px}}{x} - \frac{e^{-ux}}{e^x - 1}$ for $x > 0$, $g_{p,u}(0) := u + \frac{1}{2} - p$, $h_u(x) := \frac{e^{-ux} x}{e^x - 1}$ for $x > 0$, $h_u(0) := 1$. Then, for every fixed $u \geq 0$ the next statements hold:*

1. $g_{p,u} \in \Phi(l_2) \iff 0 \leq p \leq \sqrt{m_\infty(u)}$.
2. $-g_{p,u} \in \Phi(l_2) \iff p \geq \sqrt{M_\infty(u)} = u + 1$.
3. The inequality $\frac{d}{dt} \{t^{2\mu} S_\mu(t, u)\} \geq 0$ is valid for every $\mu > 0$ and $t > 0 \iff -\frac{d}{dx}(h_u(x)) \in \Phi(l_2)$.

The next corollary immediately follows from inequality $u^2 + u < m_\infty(u)$ and Theorem 2.11.

Corollary 2.12. *If $u \geq 0$ and $0 \leq p \leq \sqrt{u^2 + u}$, then $g_{p,u} \in \Phi(l_2)$.*

The proof of theorems of Bochner's, Hausdorff-Bernstein-Widder's and Schoenberg's can be found, for example, in [22, 23, 24, 25, 26].

3. EULER-MACLAURIN'S FORMULA. PROOFS OF THEOREMS 2.1, 2.2 AND 2.3

Lemma 3.1. 1. *Let $p \in \mathbb{N}$, $u > -p$, $\varepsilon > 0$, $n \in \mathbb{Z}_+$. If $F \in C^n[\varepsilon u^-, \varepsilon(p+u)]$, where $u^- = \frac{u-|u|}{2}$, then the next equality*

$$(3.1) \quad \sum_{k=1}^p F(\varepsilon k + \varepsilon u) = \frac{1}{\varepsilon} \int_0^{\varepsilon(p+u)} F(t) dt + \sum_{k=0}^n \frac{(-1)^{k+1} \varepsilon^k}{(k+1)!} (B_{k+1}(0) F^{(k)}(\varepsilon p + \varepsilon u) - B_{k+1}(-u) F^{(k)}(0)) + \frac{(-1)^n \varepsilon^n}{(n+1)!} \left(\int_0^{\varepsilon p} b_{n+1} \left(\frac{t}{\varepsilon} \right) dF^{(n)}(t + \varepsilon u) + \int_0^1 B_{n+1}(-u + ut) dF^{(n)}(t\varepsilon u) \right)$$

valid.

2. *Let $p \in \mathbb{N}$, $u > -2p$, $\varepsilon > 0$, $n \in \mathbb{Z}_+$. If $G \in C^n[\varepsilon u^-, \varepsilon(2p+u)]$, then the next equality*

$$(3.2) \quad \sum_{k=1}^{2p} (-1)^{k-1} G(\varepsilon k + \varepsilon u) = \sum_{k=0}^n \frac{(-1)^{k+1} \varepsilon^k}{2 k!} (E_k(0) G^{(k)}(2\varepsilon p + \varepsilon u) - E_k(-u) G^{(k)}(0)) + \frac{(-1)^n \varepsilon^n}{2 n!} \left(\int_0^{2\varepsilon p} e_n \left(\frac{t}{\varepsilon} \right) dG^{(n)}(t + \varepsilon u) + \int_0^1 E_n(-u + ut) dG^{(n)}(t\varepsilon u) \right)$$

valid. Here, $B_n(x)$ and $E_n(x)$ are Bernoulli's and Euler's polynomial, and $b_n(x)$ and $e_n(x)$ Bernoulli's and Euler's splines.

Proof. Let us consider the statement **1**. Under our conditions, the points 0 and εu belong to the segment $[\varepsilon u^-, \varepsilon(p+u)]$. Consequently, all integrals in (3.1) are well defined. Obviously, the function $b_n(x)$ are periodic of period $T = 1$. It follows from the properties of Bernoulli's polynomials that $b_0(x) = 1$, $b_1(x) = \frac{1}{2} + \{x\}$. For $n \geq 2$, the functions $b_n(x)$ are absolutely continuous on $\mathbb{R}$, and $b_n \in C^{n-2}(\mathbb{R})$, $b'_n(x) = n b_{n-1}(x)$ for $n \geq 3$, $x \in \mathbb{R}$ and $b'_2(x) = 2b_1(x)$ for $x \notin \mathbb{Z}$. Taking into account these properties and applying the integration in part in the Riemann-Stieltjes's

integrals formula, we get

$$\begin{aligned} \frac{(-1)^n \varepsilon^n}{(n+1)!} \int_0^{\varepsilon p} b_{n+1} \left(\frac{t}{\varepsilon} \right) dF^{(n)}(t + \varepsilon u) &= \frac{(-1)^n \varepsilon^n}{(n+1)!} b_{n+1}(0) (F^{(n)}(\varepsilon p + \varepsilon u) - F^{(n)}(\varepsilon u)) + \\ \frac{(-1)^{n-1} \varepsilon^{n-1}}{n!} \int_0^{\varepsilon p} b_n \left(\frac{t}{\varepsilon} \right) dF^{(n-1)}(t + \varepsilon u) &= \dots = \\ \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{(k+1)!} b_{k+1}(0) (F^{(k)}(\varepsilon p + \varepsilon u) - F^{(k)}(\varepsilon u)) &- \int_0^{\varepsilon p} F(t + \varepsilon u) db_1 \left(\frac{t}{\varepsilon} \right). \end{aligned}$$

Taking into account equality $b_1(x) = -\frac{1}{2} + x - [x]$, we obtain

$$\begin{aligned} \int_0^{\varepsilon p} F(t + \varepsilon u) db_1 \left(\frac{t}{\varepsilon} \right) &= \frac{1}{\varepsilon} \int_0^{\varepsilon p} F(t + \varepsilon u) dt - \int_0^{\varepsilon p} F(t + \varepsilon u) d \left[\frac{t}{\varepsilon} \right] = \\ &= \frac{1}{\varepsilon} \int_0^{\varepsilon p} F(t + \varepsilon u) dt - \sum_{k=1}^p F(\varepsilon k + \varepsilon u). \end{aligned}$$

Then

$$\begin{aligned} (3.3) \quad \frac{(-1)^n \varepsilon^n}{(n+1)!} \int_0^{\varepsilon p} b_{n+1} \left(\frac{t}{\varepsilon} \right) dF^{(n)}(t + \varepsilon u) &= \\ \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{(k+1)!} b_{k+1}(0) (F^{(k)}(\varepsilon p + \varepsilon u) - F^{(k)}(\varepsilon u)) &- \frac{1}{\varepsilon} \int_{\varepsilon u}^{\varepsilon(p+u)} F(t) dt + \sum_{k=1}^p F(\varepsilon k + \varepsilon u). \end{aligned}$$

By the same, taking into account the equality $B'_n(x) = nB_{n-1}(x)$, $n \in \mathbb{N}$ and applying the integration in part formula, we get the next equality

$$\begin{aligned} \frac{(-1)^n \varepsilon^n}{(n+1)!} \int_0^1 B_{n+1}(-u + ut) dF^{(n)}(t\varepsilon u) &= \\ \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{(k+1)!} (B_{k+1}(0)F^{(k)}(\varepsilon u) - B_{k+1}(-u)F^{(k)}(0)) &- \int_0^1 F(t\varepsilon u) dB_1(-u + ut). \end{aligned}$$

Taking into account equality $B_1(x) = -\frac{1}{2} + x$, we obtain

$$\begin{aligned} (3.4) \quad \frac{(-1)^n \varepsilon^n}{(n+1)!} \int_0^1 B_{n+1}(-u + ut) dF^{(n)}(t\varepsilon u) &= \\ \sum_{k=0}^n \frac{(-1)^k \varepsilon^k}{(k+1)!} (B_{k+1}(0)F^{(k)}(\varepsilon u) - B_{k+1}(-u)F^{(k)}(0)) &- \frac{1}{\varepsilon} \int_0^{\varepsilon u} F(t) dt. \end{aligned}$$

Adding (3.3) and (3.4), taking into account $b_k(0) = B_k(0)$, we get equality (3.1).

The statement **2** follows from the statement **1** and the next equality

$$(3.5) \quad \sum_{k=1}^{2p} (-1)^{k-1} G(\varepsilon k + \varepsilon u) = \sum_{k=1}^{2p} G(\varepsilon k + \varepsilon u) - 2 \sum_{k=1}^p G(2\varepsilon k + \varepsilon u)$$

by application the formula (3.1), for $F = G$, to the right hand side of (3.5): to the first term changing p to $2p$, to the second term - changing ε to 2ε and u to $\frac{u}{2}$, taking into account that $\left(\frac{u}{2}\right)^- = \frac{u^-}{2}$. $\square$

Proof of Theorem 2.1. At first we will prove the statement **1**. From Lemma 3.1, by passing to the limit as $p \rightarrow +\infty$ in (3.1), we get (2.5). From (2.5) the inequality (2.6) follows. If, in addition, $F^{(n)}$ is absolutely continuous on $[0, +\infty)$, then $F^{(n+1)} \in L[0, +\infty)$ and, consequently, $\int_0^\infty |F^{(n+1)}(t+h) - F^{(n+1)}(t)| dt = o(1)$ as $h \rightarrow +0$. Furthermore, the next equality

$$|L_n(\varepsilon, u)| \leq \left| \int_0^\infty b_{n+1} \left(\frac{t}{\varepsilon} \right) F^{(n+1)}(t) dt \right| + \sup_{0 \leq x \leq 1} |b_{n+1}(x)| \int_0^\infty |F^{(n+1)}(t + \varepsilon u) - F^{(n+1)}(t)| dt + \sup_{-u \leq x \leq 0} |B_{n+1}(x)| V_0^{\varepsilon u}(F^{(n)})$$

holds. Because $b_{n+1}(x)$ is periodic of period $T = 1$, $F^{(n+1)} \in L[0, +\infty)$, and $\int_0^1 b_{n+1}(x) dx = \int_0^1 \frac{B'_{n+2}(x)}{n+2} dx = \frac{B_{n+2}(1) - B_{n+2}(0)}{n+2} = 0$ for every $n \in \mathbb{Z}_+$, the Riemann-Lebesgue theorem imply that the first term of last inequality tends to zero as $\varepsilon \rightarrow +0$. The second term tends to zero by arguments above. Finally, $V_0^{\varepsilon u}(F^{(n)}) \rightarrow 0$ as $\varepsilon \rightarrow +0$ because $F^{(n)}$ is a bounded variation on $[0, +\infty)$ and continuous at zero function. Thus, $L_n(\varepsilon, u) = o(1)$ as $\varepsilon \rightarrow +0$, and estimation in $o(1)$ is uniform in $u \in [0, a]$ for every fixed $a > 0$.

The statement **2** can be proved in a similar way. Taking in mind that, if in addition $F^{(n)}$ is absolutely continuous on $[q, +\infty)$, $q < 0$, then $F^{(n+1)} \in L[q, +\infty)$ and, consequently, $\int_0^\infty |F^{(n+1)}(t+h) - F^{(n+1)}(t)| dt = o(1)$ for $h \rightarrow -0$ and $V_{\varepsilon u}^0(F^{(n)}) \rightarrow 0$ as $\varepsilon \rightarrow +0$ since $F^{(n)}$ is a bounded variation on $[q, +\infty)$ and continuous at zero functions. $\square$

Proof of Theorem 2.2. The proof is a similar to the proof of Theorem 2.1. Let us prove, for example, the statement **1**. The equality (2.8) follows from Lemma 3.1 by passing to the limit in (3.2) as $p \rightarrow +\infty$. The inequality (2.9) follows from (2.8). If, in addition, $G^{(n)}$ is absolutely continuous on $[0, +\infty)$, then

$$|l_n(\varepsilon, u)| \leq \left| \int_0^\infty e_n \left(\frac{t}{\varepsilon} \right) G^{(n+1)}(t) dt \right| + \sup_{0 \leq x \leq 2} |e_n(x)| \int_0^\infty |G^{(n+1)}(t + \varepsilon u) - G^{(n+1)}(t)| dt + \sup_{-u \leq x \leq 0} |E_n(x)| V_0^{\varepsilon u}(G^{(n)}).$$

It is obviously that $e_n(x)$ is periodic of period $T = 2$ function and $\int_0^2 e_n(x) dx = 0$ and so on. $\square$

Proof of Theorem 2.3. Let us prove the statement **1**. From the conditions follows the existence of finite limit $F^{(n)}(+\infty) = \int_q^\infty F^{(n+1)}(x) dx + F^{(n)}(q)$ for every $n \in \mathbb{Z}_+$. Since $F^{(n)} \in L[q, +\infty)$, than $F^{(n)}(+\infty) = 0$. Further, the Theorem 2.1 can be applied to the function F for every $n \in \mathbb{Z}_+$.

The statement **2** can be proved in a similar way. From the conditions follows the existence of finite limit $G^{(n)}(+\infty) = 0$ for every $n \in \mathbb{Z}_+$. Further, the Theorem 2.2 can be applied to the function G for every $n \in \mathbb{Z}_+$. $\square$

4. THE INEQUALITIES AND ASYMPTOTIC FOR MATHIEU'S SERIES. PROOF OF THEOREM 2.4

Lemma 4.1. *Let $\gamma \geq 0$, $\alpha > 0$, $\delta := \alpha(\mu + 1) - \gamma > 0$, $g(x) := x^\gamma(x^\alpha + 1)^{-\mu-1}$. Then*

1. $g \in C[0, +\infty) \cap C^\infty(0, +\infty)$ and $g^{(r)} \in L[1, +\infty)$, $g^{(r-1)}(+\infty) = 0$ for every $r \in \mathbb{N}$. Furthermore, $g \in L[1, +\infty) \iff \delta > 1$.
2. If $(\gamma, \alpha) \in \mathbb{Z}_+ \times \mathbb{N}$, then $g \in C^\infty(-1, +\infty)$ and for every $p \in \mathbb{Z}_+$ the equality

$$\frac{g^{(p)}(0)}{p!} = \begin{cases} \frac{(-1)^k \Gamma(\mu+k+1)}{\Gamma(\mu+1)\Gamma(k+1)} & , \quad \text{ecnu } p = k\alpha + \gamma, k \in \mathbb{Z}_+, \\ 0 & , \quad \text{ecnu } p \neq k\alpha + \gamma, k \in \mathbb{Z}_+ \end{cases}$$

valid.

3. If $(\gamma, \alpha) \notin \mathbb{Z}_+ \times \mathbb{N}$, then the function g has finite smoothness of zero

$$r = \begin{cases} [\gamma] & , \quad \text{ecnu } \gamma \notin \mathbb{Z}_+, \\ \gamma + [\alpha] & , \quad \text{ecnu } \gamma \in \mathbb{Z}_+, \alpha \notin \mathbb{Z}_+. \end{cases}$$

In this case, $g \in C^r[0, 1]$, $g \notin C^{r+1}[0, 1]$ and $g^{(r+1)} \in L(0, 1)$. If $\gamma \notin \mathbb{Z}_+$ then $g^{(p)}(0) = 0$ for every integer $p \in [0, r]$. If $\gamma \in \mathbb{Z}_+$, $\alpha \notin \mathbb{N}$ then $g^{(p)}(0) = 0$ for every integer $p \in [0, r]$, $p \neq \gamma$ and $g^{(\gamma)}(0) = \gamma!$.

Proof. Proof follows from the next equalities

$$\begin{aligned} \frac{1}{(u+1)^{\mu+1}} &= \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(\mu+k+1)}{\Gamma(\mu+1)\Gamma(k+1)} u^k, \quad |u| < 1, \\ g(x) &= \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(\mu+k+1)}{\Gamma(\mu+1)\Gamma(k+1)} \cdot x^{\gamma+k\alpha}, \quad 0 \leq x < 1, \\ g(x) &= \sum_{k=0}^{\infty} \frac{(-1)^k \Gamma(\mu+k+1)}{\Gamma(\mu+1)\Gamma(k+1)} \cdot \frac{1}{x^{\delta+k\alpha}}, \quad x > 1. \end{aligned} \tag{4.1}$$

□

Proof of Theorem 2.4. follows from Lemma 4.1, the Theorems 2.1 and 2.2 for $G = F = g$ and $\varepsilon = \frac{1}{t}$. □

Corollary 4.2. Let $\gamma \geq 0$, $\alpha > 0$, $\delta := \alpha(\mu+1) - \gamma > 0$ and $g(x) := x^\gamma(x^\alpha+1)^{-\mu-1}$. Then, on the interval $[0, +\infty)$, the g is a bounded variation function which equal to $V_0^\infty(g) = 1$ if $\gamma = 0$ and $V_0^\infty(g) = 2 \left(\frac{\gamma}{\delta}\right)^{\frac{2}{\alpha}} \left(\frac{\gamma}{\delta} + 1\right)^{-\mu-1}$ if $\gamma > 0$. Furthermore, the equality

$$\tilde{S}(t, u, \gamma, \alpha, \mu) = \frac{g(0) + l(t, u)}{t^{\alpha(\mu+1)-\gamma}}, \quad \text{where } |l(t, u)| \leq V_0^\infty(g), \quad l(t, u) = o(1), \quad t \rightarrow +\infty, \tag{4.2}$$

holds for every $u \geq 0$ and $t > 0$. If, in addition, $\delta > 1$, then the equality

$$S(t, u, \gamma, \alpha, \mu) = \frac{\frac{2}{\alpha} B\left(\frac{\gamma+1}{\alpha}, \mu+1 - \frac{\gamma+1}{\alpha}\right)}{t^{\alpha(\mu+1)-\gamma-1}} - \frac{(1+2u)g(0) - L(t, u)}{t^{\alpha(\mu+1)-\gamma}} \tag{4.3}$$

holds for every $u \geq 0$ and $t > 0$, where $|L(t, u)| \leq (1+2u)V_0^\infty(g)$ and $L(t, u) = o(1)$ as $t \rightarrow +\infty$. In both cases, the estimation in $o(1)$ is uniform in $u \in [0, a]$ for every fixed $a > 0$.

Proof. In the case $\gamma = 0$, the function g strictly decrease on $[0, +\infty)$ and, consequently, $V_0^\infty(g) = g(0) - g(+\infty) = 1$. In the case $\gamma > 0$, the function g increases on $[0, x_0]$ and decreases on $[x_0, +\infty)$, where $x_0 = \left(\frac{\gamma}{\delta}\right)^{1/\alpha}$ and, consequently, $V_0^\infty(g) = 2g(x_0)$. Further, we apply the statements 1 of Theorems 2.2 and 2.1 with $n = 0$, $F = G = g$, and $\varepsilon = \frac{1}{t}$. The inequalities (2.8) and (2.5) with $n = 0$ yields $l(t, u) = l_0\left(\frac{1}{\varepsilon}, u\right)$ and $L(t, u) = 2L_0\left(\frac{1}{\varepsilon}, u\right)$. Then we use the inequalities (2.9) and

(2.6), having in mind that $B_1(x) = -\frac{1}{2} + x$, $b_1(x) = -\frac{1}{2} + \{x\}$, $E_0(x) = 1$, $e_0(x) = 1$ for $0 \leq x < 1$, and $e_0(x) = -1$ for $1 \leq x < 2$, and $V_0^{\varepsilon u}(g) + V_{\varepsilon u}^{\infty}(g) = V_0^{\infty}(g)$. $\square$

Corollary 4.3. *Let $(\frac{\gamma}{2}, \frac{\alpha}{2}) \in \mathbb{Z}_+ \times \mathbb{N}$, $\delta := \alpha(\mu+1) - \gamma > 0$. Then, for every $n \in \mathbb{N}$ the equality*

$$(4.4) \quad \tilde{S}(t, 0, \gamma, \alpha, \mu) = \frac{(-1)^\gamma E_\gamma(0)}{t^{\alpha(\mu+1)}} + o\left(\frac{1}{t^{\alpha(n+\mu+1)}}\right), \quad t \rightarrow +\infty$$

holds. If, in addition $\delta > 1$, then, for every $n \in \mathbb{N}$, the equality

$$(4.5) \quad S(t, 0, \gamma, \alpha, \mu) = \frac{1}{t^{\alpha(\mu+1)-\gamma-1}} \cdot \frac{2}{\alpha} B\left(\frac{\gamma+1}{\alpha}, \mu+1 - \frac{\gamma+1}{\alpha}\right) + \frac{(-1)^\gamma 2 B_{\gamma+1}(0)}{(\gamma+1) t^{\alpha(\mu+1)}} + o\left(\frac{1}{t^{\alpha(n+\mu+1)}}\right), \quad t \rightarrow +\infty$$

holds.

Proof. Proof immediately follows from Theorems 2.1, 2.2 and 2.4, taking into account, that $E_m(0) = \frac{2(1-2^{m+1})}{m+1} B_{m+1}(0)$, for $m \in \mathbb{Z}_+$, $B_1(0) = -\frac{1}{2}$, and $B_{2p+1}(0) = 0$, for $p \in \mathbb{N}$. Thus, $E_0(0) = 1$ and $E_{2p}(0) = 0$, for $p \in \mathbb{N}$. $\square$

Remark 4.1. It seems to be remainder term in the Corollary 4.3 approaching zero more foster. It can be seen in the next examples. With using the summation Poisson's formula, we get (see, for example, [27, Example 11.3]),

$$\sum_{k=1}^{\infty} \frac{2}{k^2 + t^2} = \frac{\pi}{t} - \frac{1}{t^2} + \frac{2\pi}{t(e^{2\pi t} - 1)} = \frac{\pi}{t} - \frac{1}{t^2} + O\left(\frac{1}{te^{2\pi t}}\right), \quad t \rightarrow +\infty.$$

From this and the equality (2.3) follows that

$$\sum_{k=1}^{\infty} \frac{2(-1)^{k-1}}{k^2 + t^2} = \frac{1}{t^2} - \frac{2\pi}{t(e^{\pi t} - e^{-\pi t})} = \frac{1}{t^2} + O\left(\frac{1}{te^{\pi t}}\right), \quad t \rightarrow +\infty.$$

The last example (see, for example, [28, §3.11–3.12]):

$$\sum_{k=1}^{\infty} \frac{2(-1)^{k-1}}{\sqrt{k^2 + t^2}} = \frac{1}{t} + O\left(\frac{1}{\sqrt{t}e^{\pi t}}\right), \quad t \rightarrow +\infty.$$

5. HERMITE-HADAMARD'S INEQUALITY. PROOF OF THEOREMS 2.5 AND 2.6

Proof of Theorem 2.5. Let $d \geq 0$, $\beta > 0$. Then

$$S(t) \leq (\geq) \frac{C}{(t^\beta + d)^{\frac{\delta_1}{\beta}}}, \quad t > 0 \iff d \leq (\geq) f_\beta(t) := \left(\frac{C}{S(t)}\right)^{\frac{\beta}{\delta_1}} - t^\beta, \quad t > 0.$$

It is obviously, at $t \rightarrow +\infty$, the relations

$$f_\beta(t) = t^\beta \left(\left(\frac{C}{S(t)t^{\delta_1}} \right)^{\frac{\beta}{\delta_1}} - 1 \right) \sim \frac{t^\beta \beta}{\delta_1} \left(\frac{C}{S(t)t^{\delta_1}} - 1 \right) = \frac{t^\beta \beta}{\delta_1} \frac{C - S(t)t^{\delta_1}}{S(t)t^{\delta_1}} \sim \frac{t^{\beta-\beta_1} \beta A}{\delta_1 C}$$

holds. Therefore, if $\beta < \beta_1$, the inequality $0 < d \leq f_\beta(t)$, is impossible for big t (otherwise $0 < d \leq 0$). If $\beta > \beta_1$, the inequality $d \geq f_\beta(t)$ is impossible for big t (otherwise $d \geq +\infty$).

Let $\beta = \beta_1$. Then $f(t) = f_{\beta_1}(t) > 0$ for $t \geq 0$, and $f(+\infty) = \frac{\beta_1 A}{\delta_1 C} > 0$. Therefore, $m = \inf_{t \geq 0} f(t) > 0$, $M = \sup_{t \geq 0} f(t) < +\infty$. Then, for $d \geq 0$, the next statements

$$S(t) \leq (\geq) \frac{C}{(t^{\beta_1} + d)^{\frac{\delta_1}{\beta_1}}}, \quad t > 0 \iff d \leq (\geq) f(t), \quad t > 0 \iff d \leq m \ (d \geq M)$$

are obvious. $\square$

Lemma 5.1. *Let g is convex downwards on the interval I function, that is, for every $x, y \in I$, and $\lambda \in (0, 1)$, the inequality $g(\lambda x + (1 - \lambda)y) \leq \lambda g(x) + (1 - \lambda)g(y)$ holds. Then, for every $a, b \in I : a < b$, the Hermite-Hadamard's inequality*

$$(5.1) \quad (b - a) g\left(\frac{a + b}{2}\right) \leq \int_a^b g(x) dx \leq (b - a) \frac{g(a) + g(b)}{2}$$

valid. The equality in one inequality of (5.1) occurs iff g is linear on the segment $[a, b]$.

Proof. Note, that every convex downwards on the interval I function f is continuous on that interval and, for every points $x_1, x_2, x_3 \in I : x_1 < x_2 < x_3$ the inequalities

$$(5.2) \quad \frac{f(x_2) - f(x_1)}{x_2 - x_1} \leq \frac{f(x_3) - f(x_1)}{x_3 - x_1} \leq \frac{f(x_3) - f(x_2)}{x_3 - x_2}$$

hold. Therefore, $g \in C(I)$. The function $h(x) := g\left(\frac{a+b}{2} + x\right) + g\left(\frac{a+b}{2} - x\right)$ is convex downwards on $[-c, c]$ with $c = \frac{b-a}{2} > 0$ and, furthermore, is even function. Therefore, h is increases on $[0, c]$ (it is consequence of (5.2) with $f = h$, $-x_1 = x_3 \in (0, c]$, and $x_2 \in (0, x_3)$) and, therefore, the inequalities

$$\int_a^b g(x) dx = \int_0^c h(x) dx \leq c h(c) (\geq c h(0))$$

hold. The inequality (5.1) proved. If g is a linear on $[a, b]$, then both inequalities of (5.1) reduces to the equalities and conversely. If one inequality of (5.1) becomes equality then h is a constant on $[-c, c]$. Therefore, the function g is a convex downwards and upwards simultaneously on $[a, b]$. In this case, for $f = g$, the inequalities in (5.2) reduces to the equalities with $x_1 = a$, $x_3 = b$ and, consequently, g is a linear function on the segment $[a, b]$. $\square$

Lemma 5.2. *Let g is convex downwards on $(0, +\infty)$ function such that the integral $\int_1^{+\infty} g(x) dx$ is convergent. Then g decreases on $(0, +\infty)$, $g(+\infty) = 0$ and the next statements hold:*

1. *For every $u \geq -1$, $y > -u^2 - u$ the inequality*

$$(5.3) \quad \sum_{k=1}^{\infty} 2(k + u) g((k + u)^2 + y) \leq \int_{u^2 + u + y}^{+\infty} g(x) dx$$

holds. The equality in (5.3) occurs iff the function g is a linear on each segment $[a_k, a_{k+1}]$, $k \in \mathbb{N}$, with $a_k = (k - \frac{1}{2} + u)^2 + y - \frac{1}{4}$.

2. *For every $u \geq -\frac{3}{2}$, $y > -(1 + u)^2$ the inequality*

$$(5.4) \quad \int_{(1+u)^2 + y}^{+\infty} g(x) dx + \left(\frac{1}{2} + u\right) g((1 + u)^2 + y) \leq \sum_{k=1}^{\infty} 2(k + u) g((k + u)^2 + y)$$

holds. The equality in (5.4) occurs iff the function g is a linear on each segment $[s_k, s_{k+1}]$, $k \in \mathbb{N}$, with $s_k = (k + u)^2 + y$.

Proof. The inequality (5.1) implies

$$\frac{2}{t-a} \int_a^t g(x) dx - g(a) \leq g(t) \leq \frac{1}{2(t-a)} \int_a^{2t-a} g(x) dx, \quad t > a > 0.$$

Therefore, $\lim_{t \rightarrow +\infty} \frac{g(t)}{t} = 0$. Passing to the limit, as $x_3 \rightarrow +\infty$, in the inequality (5.2) with $f = g$, we conclude that g decreases on $(0, +\infty)$. The monotonicity of g and convergence of the integral $\int_1^{+\infty} g(x) dx$ imply $g(+\infty) = 0$.

Let us prove the statement **1**. In the left inequality of (5.1), we can take $a = a_k$ and $b = a_{k+1}$, since $a_{k+1} - a_k = 2(k+u) \geq 0$ for every $k \in \mathbb{N}$, and $a_1 = u^2 + u + y > 0$. Then $\frac{a+b}{2} = (k+u)^2 + y$. By summation of these inequalities with respect to $k \in \mathbb{N}$, we get the inequality (5.3). In this case, the equality in (5.3) occurs (Lemma 5.1) only if g is a linear function on each segment $[a_k, a_{k+1}]$, $k \in \mathbb{N}$.

Let us prove the statement **2**. In the right inequality of (5.1), we can take $a = s_k$ and $b = s_{k+1}$, since $s_{k+1} - s_k = 2(k+u) + 1 \geq 0$, for every $k \in \mathbb{N}$, and $s_1 = (1+u)^2 + y > 0$. Then

$$\begin{aligned} \int_{s_k}^{s_{k+1}} g(x) dx &\leq \left(k+u + \frac{1}{2}\right) (g((k+u)^2 + y) + g((k+1+u)^2 + y)) = \\ &= (k+u)g((k+u)^2 + y) + (k+1+u)g((k+1+u)^2 + y) + \\ &\quad \frac{1}{2}(g((k+u)^2 + y) - g((k+1+u)^2 + y)). \end{aligned}$$

We add these inequalities with respect to $k \in \mathbb{N}$. By the inequality (5.3), in the right hand side, we get absolutely convergent series

$$\int_{s_1}^{+\infty} g(x) dx \leq \sum_{k=1}^{\infty} 2(k+u)g((k+u)^2 + y) - \left(\frac{1}{2} + u\right) g((1+u)^2 + y).$$

The last inequality is equivalent to the inequality (5.4), and the equality occurs (see Lemma 5.1) only if g is a linear function on each segment $[s_k, s_{k+1}]$, $k \in \mathbb{N}$. $\square$

Proof of Theorem 2.6. Let us prove the property **1**. The series (2.18) converge uniformly on each compact $K \subset E$ and, consequently, $S(u, y) \in C(E)$. In fact, for every compact $K \subset E$, the number $n_0 = n_0(K)$ does exist such that, for every number $n \geq n_0$ and every point $(u, y) \in K$ the inequalities $n+u \geq -1$ and $(n+u)^2 + (n+u) + y \geq n$ hold. Hence, the inequality (5.3) can be applied to the remainder of the series (2.18):

$$\begin{aligned} \sum_{k=n+1}^{\infty} 2(k+u)g((k+u)^2 + y) &= \sum_{k=1}^{\infty} 2(k+n+u)g((k+n+u)^2 + y) \leq \\ &\int_{(n+u)^2 + (n+u) + y}^{+\infty} g(x) dx \leq \int_n^{+\infty} g(x) dx, \quad n \geq n_0, (u, y) \in K. \end{aligned}$$

From the last inequality and nonnegativity of g , the uniform convergence of series (2.18) on K follows.

If $(u_0, y_0) \in E$ the $(u_0, y) \in E$ for every $y > y_0$ and $S(u_0, +\infty) = 0$. Each term of the series (2.18) and every their remainder series with the number, satisfying to the inequalities (see above) $n+u_0 \geq -1$ and $(n+u_0)^2 + (n+u_0) + y_0 > 0$, obeys this property.

For every fixed $u \geq -1$, the function $S(u, y)$ is nonnegative, decreasing, and convex downwards with respect to $y \in (-(1+u)^2, +\infty)$, since each term of the series (2.18) obeys this property.

If, in addition, $g(x) > 0$ for every $x > 0$ then in the inequality (5.2) for $f = g$, we can always choose $x_3 > x_2$ such that the inequality $f(x_3) < f(x_2)$ holds. Therefore, the function g strictly decreases on $(0, +\infty)$ and, consequently, the function $S(u, y)$ strictly decreases with respect to $y \in (-(1+u)^2, +\infty)$ for every fixed $u \geq -1$.

Note that, if for some $u \geq -1$ and $y_0 > -(1+u)^2$, $S(u, y) \equiv \text{const}$ for $y \geq y_0$, then $S(u, y) \equiv 0$, for $y \geq y_0$, and, consequently, $g(x) \equiv 0$ for $x \geq x_0$, where $x_0 = (1+u)^2 + y_0$ if $u > -1$ and $x_0 = 1 + y_0$ if $u = -1$.

The property **2** includes in the Lemma 5.2. The Theorem 2.6 is proved. $\square$

6. $S_\mu(t, u)$ SERIES AND POSITIVE DEFINITENESS. PROOFS OF THEOREMS 2.7, 2.8, 2.9 AND 2.11

Lemma 6.1. 1. *For every $(p, t) \in \mathbb{R}^2 \setminus \{(0, s) : s \leq 0\}$, $\nu > \mu > 0$, and $u \in \mathbb{R}$, the inequality*

$$(6.1) \quad \int_t^{+\infty} \left(\frac{1}{\nu(p^2 + y^2)^\nu} - S_\nu(y, u) \right) y |y^2 - t^2|^{\nu-\mu-1} dy = \frac{B(\nu - \mu, \mu + 1)}{2} \left(\frac{1}{\mu(p^2 + t^2)^\mu} - S_\mu(t, u) \right)$$

holds.

2. *For every $u, t, b \in \mathbb{R}$, $\nu > \mu > 0$, the inequality*

$$(6.2) \quad \int_t^{+\infty} (\nu S_\nu(y, u) - (\nu + 1) S_{\nu+1}(y, u)(y^2 + b)) y |y^2 - t^2|^{\nu-\mu-1} dy = \frac{B(\nu - \mu, \mu + 1)}{2} (\mu S_\mu(t, u) - (\mu + 1) S_{\mu+1}(t, u)(t^2 + b))$$

holds.

Proof. At first, we prove that if the condition

1) $(p, t) \in \mathbb{R}^2 \setminus \{(0, s) : s \leq 0\}$, $\nu > \mu > 0$ or 2) $p = 0$, $t < 0$, $1 > \nu > \mu > 0$

holds, then the equality

$$(6.3) \quad \int_t^{+\infty} \frac{y |y^2 - t^2|^{\nu-\mu-1}}{(p^2 + y^2)^\nu} dy = \frac{B(\nu - \mu, \mu)}{2} \cdot \frac{1}{(p^2 + t^2)^\mu}$$

take place. It is easy check that the integral in the left hand side of (6.3) converges absolutely if the condition 1) or 2) holds. In any case, by substitution of variable $s = \frac{y^2 - t^2}{p^2 + t^2}$ in (6.3), we get

$$\frac{1}{2} \cdot \frac{1}{(p^2 + t^2)^\mu} \int_0^{+\infty} \frac{s^{\nu-\mu-1}}{(1+s)^\nu} ds = \frac{B(\nu - \mu, \mu)}{2} \cdot \frac{1}{(p^2 + t^2)^\mu}.$$

The equality (6.3) proved. After substitution in (6.3) ν by $\nu + 1$ and μ by $\mu + 1$, the equality

$$\int_t^{+\infty} \frac{y |y^2 - t^2|^{\nu-\mu-1}}{(p^2 + y^2)^{\nu+1}} dy = \frac{B(\nu - \mu, \mu + 1)}{2} \cdot \frac{1}{(p^2 + t^2)^{\mu+1}}$$

take place for every $(p, t) \in \mathbb{R}^2 \setminus \{(0, s) : s \leq 0\}$, $\nu > \mu > -1$. Tacking into account the last equality, for every $u, t \in \mathbb{R}$, $\nu > \mu > 0$, we get

$$(6.4) \quad \int_t^{+\infty} S_\nu(y, u) y |y^2 - t^2|^{\nu-\mu-1} dy = \sum_{k=1}^{\infty} 2(k+u) \int_t^{+\infty} \frac{y |y^2 - t^2|^{\nu-\mu-1}}{((k+u)^2 + y^2)^{\nu+1}} dy = \frac{B(\nu-\mu, \mu+1)}{2} S_\mu(t, u).$$

Note that, in the series (2.30) for $S_\nu(y, u)$ and $S_\mu(t, u)$, the terms for $k = -u$ we absent if $-u \in \mathbb{N}$.

Taking into account the equality $\frac{B(\nu-\mu, \mu)}{\nu} = \frac{B(\nu-\mu, \mu+1)}{\mu}$, from (6.3) and (6.4) the equality (6.1) follows.

The integral in the left hand side of (6.2) is equal $I_1 - I_2 - I_3(t^2 + b)$, where

$$\begin{aligned} I_1 &= \nu \int_t^{+\infty} S_\nu(y, u) y |y^2 - t^2|^{\nu-\mu-1} dy = \frac{B(\nu-\mu, \mu+1)}{2} \nu S_\mu(t, u), \\ I_2 &= (\nu+1) \int_t^{+\infty} S_{\nu+1}(y, u) y |y^2 - t^2|^{\nu-\mu} dy = \frac{B(\nu-\mu, \mu+1)}{2} (\nu-\mu) S_\mu(t, u), \\ I_3 &= (\nu+1) \int_t^{+\infty} S_{\nu+1}(y, u) y |y^2 - t^2|^{\nu-\mu-1} dy = \frac{B(\nu-\mu, \mu+1)}{2} (\mu+1) S_{\mu+1}(t, u). \end{aligned}$$

To calculate I_1 we used (6.4). By changing ν by $\nu+1$ in (6.4) we calculated I_2 . To calculate I_3 , we changed ν by $\nu+1$, and μ by $\mu+1$ in (6.4). The equality (6.2) proved. $\square$

For functions h defined on $(0, +\infty)$, let us define a Hankel transform¹ for $m \in \mathbb{C}$ and $t > 0$:

$$(6.5) \quad \mathfrak{F}_m(h)(t) := t^{1-\frac{m}{2}} \int_0^{+\infty} h(x) x^{\frac{m}{2}} J_{\frac{m}{2}-1}(tx) dx = \int_0^{+\infty} h(x) x^{m-1} j_{\frac{m}{2}-1}(tx) dx,$$

where J_λ is a Bessel function of the first kind, and

$$(6.6) \quad j_\lambda(x) := \frac{J_\lambda(x)}{x^\lambda} = \frac{1}{2^\lambda} \sum_{k=0}^{\infty} \frac{1}{\Gamma(k+\lambda+1)} \cdot \frac{\left(-\frac{x^2}{4}\right)^k}{k!}, \quad x \in \mathbb{C}, \quad \lambda \in \mathbb{C}.$$

For $m \in \mathbb{N}$ the transform $\mathfrak{F}_m$ is related to a Fourier transform for radial functions:

$$(6.7) \quad F_m(h(\|\cdot\|_2))(x) = (2\pi)^{\frac{m}{2}} \mathfrak{F}_m(h)(\|x\|_2), \quad x \in \mathbb{R}^m.$$

Lemma 6.2. *Let*

$$(6.8) \quad \begin{aligned} g_{p,u}(x) &:= \frac{e^{-px}}{x} - \frac{e^{-ux}}{e^x - 1}, \quad g_{p,u}(0) := u + \frac{1}{2} - p; \\ G_{p,u,\mu}(x) &:= x g'_{p,u}(x) + (2\mu - 1) g_{p,u}(x); \\ h_u(x) &:= \frac{e^{-ux} x}{e^x - 1}, \quad h_u(0) := 1. \end{aligned}$$

Then equalities

$$(6.9) \quad \frac{1}{\mu(p^2 + t^2)^\mu} - S_\mu(t, u) = \frac{\sqrt{\pi} \mathfrak{F}_{2\mu+1}(g_{p,u})(t)}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)}, \quad t \geq 0, \quad p > 0, \quad u > -1, \quad \mu > 0,$$

¹A non-standard notation is used here.

$$(6.10) \quad \frac{1}{\mu(p^2 + t^2)^\mu} - S_\mu(t, u) = \frac{\sqrt{\pi} \mathfrak{F}_{2\mu-1}(G_{p,u,\mu})(t)}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1) t^2}, \quad t > 0, p > 0, u > -1, \mu > \frac{1}{2},$$

$$(6.11) \quad \frac{d}{dt} \{t^{2\mu} S_\mu(t, u)\} = -\frac{\sqrt{\pi} t^{2\mu-1} \mathfrak{F}_{2\mu+1}(h'_u)(t)}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)}, \quad t > 0, u > -1, \mu > 0$$

hold.

Proof. By [29, Chapter 4.14(7,8), $\nu = \mu - \frac{1}{2}$] we get the equalities

$$(6.12) \quad \frac{1}{\mu(p^2 + a^2)^\mu} = \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)} \int_0^{+\infty} e^{-px} x^{2\mu-1} j_{\mu-\frac{1}{2}}(ax) dx, \quad a \in \mathbb{R}, p > 0, \mu > 0,$$

$$(6.13) \quad \frac{2p}{(p^2 + a^2)^{\mu+1}} = \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)} \int_0^{+\infty} e^{-px} x^{2\mu} j_{\mu-\frac{1}{2}}(ax) dx, \quad a \in \mathbb{R}, p > 0, \mu > -\frac{1}{2}.$$

After the summation of the equalities (6.13) for $p = k + u$, $k \in \mathbb{N}$, $u > -1$, $a = t \geq 0$, $\mu > 0$, we get

$$(6.14) \quad S_\mu(t, u) = \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)} \int_0^{+\infty} \frac{e^{-ux}}{e^x - 1} x^{2\mu} j_{\mu-\frac{1}{2}}(tx) dx, \quad t \geq 0, u > -1, \mu > 0.$$

For $a = t \geq 0$, the equalities (6.14) and (6.12) imply the equality (6.9). If, in addition, $\mu > \frac{1}{2}$, then for $t > 0$ we can use integration in part in the right hand side of (6.9). Taking into account $\frac{d}{dx}(j_\lambda(tx)) = -t^2 x j_{\lambda+1}(tx)$, we get

$$\mathfrak{F}_{2\mu+1}(g_{p,u})(t) = -\frac{1}{t^2} \int_0^{+\infty} g_{p,u}(x) x^{2\mu-1} dj_{\mu-\frac{3}{2}}(tx) = \frac{1}{t^2} \int_0^{+\infty} (g_{p,u}(x) x^{2\mu-1})' j_{\mu-\frac{3}{2}}(tx) dx,$$

from which (6.10) follows.

Let $h'_u(x) := \frac{d}{dx}(h_u(x))$. If $t > 0$, by the substitution $s = tx$, in the integral from (6.14), for $u > -1$, $\mu > 0$, we get

$$\frac{d}{dt} \{t^{2\mu} S_\mu(t, u)\} = \frac{\sqrt{\pi}}{2^{\mu-\frac{1}{2}} \Gamma(\mu+1)} \int_0^{+\infty} h'_u\left(\frac{s}{t}\right) \left(-\frac{s}{t^2}\right) s^{2\mu-1} j_{\mu-\frac{1}{2}}(s) ds.$$

The last equality is equivalent to (6.11). $\square$

Lemma 6.3. *Let $p, u \in \mathbb{R}$. Then, for $g_{p,u}(x) := \frac{e^{-px}}{x} - \frac{e^{-ux}}{e^x - 1}$, $x > 0$, the next statements hold:*

1. *If $p - u \leq \frac{1}{2}$ then $g_{p,u}(x) > 0$, for every $x > 0$.*
2. *If $p - u \geq 1$ then $g_{p,u}(x) < 0$, for every $x > 0$.*
3. *If $\frac{1}{2} < p - u < 1$ then, on an interval $(0, +\infty)$, the function $g_{p,u}(x)$ is not of constant signs.*

Proof. From the equality $g_{p,u}(x) = \frac{e^{-ux}}{x(e^x - 1)} f_{p-u}(x)$, where $f_a(x) = e^{-ax}(e^x - 1) - x$, follows that we have to find such a values of $a \in \mathbb{R}$, that $f_a(x)$ preserves sign over the interval $(0, +\infty)$. It is obvious that $f_a(0) = 0$, $f'_a(x) = (1-a)e^{(1-a)x} + ae^{-ax} - 1$, $f'_a(0) = 0$, and $f''_a(x) = ((1-a)^2 e^x - a^2) e^{-ax}$.

If $0 \leq a \leq \frac{1}{2}$, $x > 0$, then $e^x > 1 \geq \frac{a^2}{(1-a)^2}$. Consequently, $f''_a(x) > 0$ and $f_a(x) > 0$ for every $x > 0$. If $a < 0$, $x > 0$, then $f_a(x) > e^x - 1 - x > 0$.

If $a = 1$, then $f_1''(x) = -e^{-x} < 0$ and, consequently, $f_1(x) < 0$ for every $x > 0$. If $a > 1$, then $f_a(x) < f_1(x) < 0$ for every $x > 0$.

If $\frac{1}{2} < a < 1$, then $f_a(+\infty) = +\infty$ and $f_a''(0) = 1 - 2a < 0$. Consequently, $f_a(x)$ does not preserve a sign on $(0, +\infty)$. $\square$

Lemma 6.4. **1.** *If for some $\nu > 0$, $b \geq 0$, $u \in \mathbb{R}$, the function $(t^2 + b)^\nu S_\nu(t, u)$ increases (decreases) with respect to $t \in [0, +\infty)$, then, for every $0 < \mu \leq \nu$ the function $(t^2 + b)^\mu S_\mu(t, u)$ strictly increases (strictly decreases) with respect to $t \in [0, +\infty)$.*

2. *If for some $\nu > 0$, $b \geq 0$, $u \geq -1$, the function $(t^2 + b)^\nu S_\nu(t, u)$ increases with respect to $t \in [0, +\infty)$, then, for every $0 < \mu \leq \nu$, $0 \leq a \leq b$ the function $(t^2 + a)^\mu S_\mu(t, u)$ strictly increases with respect to $t \in [0, +\infty)$ and, for $t \geq 0$, the inequality $\mu(t^2 + a)^\mu S_\mu(t, u) < 1$ take place. If, in addition, $(\mu, a) \neq (\nu, b)$ then, for $t > 0$, the inequality $\frac{d}{dt}((t^2 + a)^\mu S_\mu(t, u)) > 0$ take place.*

Proof. The first statement we will prove by contradiction. Let us suppose that the function $(t^2 + b)^\nu S_\nu(t, u)$ is monotonic with respect to $t \in [0, +\infty)$ but not strictly monotonic on this interval. Then

$$(6.15) \quad \begin{aligned} \frac{d}{dt}((t^2 + b)^\nu S_\nu(t, u)) &= 2t(t^2 + b)^{\nu-1} H_{\nu,b}(t, u), \quad t > 0, \quad \text{where} \\ H_{\nu,b}(t, u) &= \nu S_\nu(t, u) - (\nu + 1) S_{\nu+1}(t, u)(t^2 + b), \quad t \in \mathbb{R}, \end{aligned}$$

preserve sign for $t > 0$ and vanishing on some interval. The function $H_{\nu,b}(t, u)$ is analytic, for every fixed $u \in \mathbb{R}$, in the neighborhood of any real point, the, by the uniqueness theorem for analytic functions, $H_{\nu,b}(t, u) = 0$ for every $t \in \mathbb{R}$. The asymptotic representation (2.31) implies that

$$(6.16) \quad H_{\nu,b}(t, u) \sim \sum_{k=1}^{\infty} \frac{B_{2k}(-u) - b B_{2k-2}(-u)}{t^{2(k+\nu)}} \cdot \frac{(-1)^{k-1} \Gamma(\nu + k)}{\Gamma(\nu + 1) \Gamma(k)}, \quad t \rightarrow +\infty,$$

and, consequently, $B_{2k}(-u) - b B_{2k-2}(-u) = 0$ for every $k \in \mathbb{N}$. It is easy to check that the system of equations has no solutions (it is enough to check it for $k = 1, 2, 3$). Statement **1**, for $\mu = \nu$ proved. If $0 < \mu < \nu$, then the equality (6.2) implies that the sign of $H_{\mu,b}(t, u)$ is equal to the sign of $H_{\nu,b}(t, u)$ for $t > 0$.

Let us prove the statement **2**. Let us suppose that the function $(t^2 + b)^\nu S_\nu(t, u)$ increasing with respect to $t \in [0, +\infty)$ for some $\nu > 0$, $b \geq 0$, $u \geq -1$. From what we already proved, this function increases with respect to $t \in [0, +\infty)$ and, consequently, $H_{\nu,b}(t, u) \geq 0$ for $t > 0$. The condition $u \geq -1$ implies that $S_\mu(t, u) > 0$ for every $\mu > 0$, $t \in \mathbb{R}$. Consequently, the inequality

$$H_{\nu,a}(t, u) = H_{\nu,b}(t, u) + (b - a)(\nu + 1) S_{\nu+1}(t, u) \geq 0$$

holds for every $0 \leq a \leq b$ and $t > 0$. If, in addition, $a < b$ then $H_{\nu,a}(t, u) > 0$ for $t > 0$. If $0 < \mu < \nu$, $0 \leq a \leq b$ then the equality (6.2) implies that $H_{\mu,a}(t, u) > 0$ for $t > 0$.

Thus, for every $0 < \mu \leq \nu$, $0 \leq a \leq b$ the function $(t^2 + a)^\mu S_\mu(t, u)$ is strictly increasing with respect to $t \in [0, +\infty)$ and, consequently, the inequality

$$\mu(t^2 + a)^\mu S_\mu(t, u) < \mu \lim_{x \rightarrow +\infty} (x^2 + a)^\mu S_\mu(x, u) = 1$$

holds for $t \geq 0$. The statement **2** proved. $\square$

Proof of Theorem 2.7. From Poisson's representation (see, for example, [19, §7.3])

$$j_\lambda(t) = \frac{1}{2^\lambda \Gamma(\lambda + \frac{1}{2}) \sqrt{\pi}} \int_{-1}^1 e^{-itu} (1-u^2)^{\lambda-\frac{1}{2}} du, \quad \lambda > -\frac{1}{2}, \quad t \in \mathbb{R}$$

follows the inequality $|j_\lambda(t)| < j_\lambda(0)$, $\lambda > -\frac{1}{2}$, $t > 0$. Further, for $t \geq 0$, $p > 0$, $u > -1$, $\mu > 0$, we use the equality (6.9) and the fact that, in the cases $p-u \leq \frac{1}{2}$ or $p-u \geq 1$, by the Lemma 6.3, the function $g_{p,u}(x)$ preserves sign for $x > 0$. In these cases, for every $t > 0$, the inequality $|\mathfrak{F}_{2\mu+1}(g_{p,u})(t)| < |\mathfrak{F}_{2\mu+1}(g_{p,u})(0)|$ holds and is equivalent to the inequality (2.32). The Theorem 2.7 proved. $\square$

Proof of Theorem 2.8. The relations (2.31) and inequality (2.29) imply that, for every fixed $\mu > 0$ and $u \geq 0$, the function $S(t) = S_\mu(t, u)$, $t \geq 0$, satisfies to the conditions of the Theorem 2.5 for $C = \frac{1}{\mu}$, $\delta_1 = 2\mu$, $A = u^2 + u + \frac{1}{6}$, $\beta_1 = 2$. From the Theorem 2.5 (statement 3) follows that $f_{\mu,u}(+\infty) = u^2 + u + \frac{1}{6}$ and, for mentioned a and b the inequality (2.33) take place.

Let us prove the statement 1. The inequalities $u^2 + u < m_\mu(u) \leq M_\mu(u) < (1+u)^2$, for $\mu > 0$, $u \geq 0$, follows from the inequality (2.23) and Remark 2.2 (the cases 1 and 2) for the function $g(x) = x^{-\mu-1}$, taking into account that $S_\mu(t, u) = S(u, t^2)$, $f_{\mu,u}(t) = \psi(u, t^2)$, $m_\mu(u) = m(u)$ и $M_\mu(u) = M(u)$ (see (2.26)). To prove the inequality $u^2 + u + \frac{1}{4} < M_\mu(u)$ we use the inequality (6.9) and the Theorem from [30]. Let us define, for $m \in \mathbb{R}$,

$$(6.17) \quad \begin{aligned} H_m &:= L((0, 1), x^{m-1} dx) \cap L((1, +\infty), x^{\frac{m-1}{2}} dx), \\ H_m^1 &:= L((0, +\infty), x^{m-1} dx), \quad H_m^2 := L((0, +\infty), x^{\frac{m-1}{2}} dx). \end{aligned}$$

Theorem 6.5 (Zastavnyi [30]). **1.** Let $m > 0$ and $f \in (H_m^1 \cup H_m^2) \cap H_m$. If, for some fixed $\delta > 0$ and $\varepsilon \in \mathbb{C}$, the function f is bounded on $(0, \delta)$ and $|\mathfrak{F}_m(f)(t)| - \varepsilon \mathfrak{F}_m(f)(t) \in H_m^1$, then $\mathfrak{F}_m(f) \in H_m^1$.
2. Let $m > 0$ and $f \in (H_m^1 \cup H_m^2) \cap H_m$. If $\mathfrak{F}_m(f)(t)$ preserves sign for $t > 0$ and f is continuous at zero then $\mathfrak{F}_m(f) \in H_m^1$. If, in addition, $f(0) = 0$ then $f(x) = 0$ for almost all $x > 0$.

Let $\mu > 0$, $u \geq 0$, and $p = u + \frac{1}{2}$. Let us suppose that $\mathfrak{F}_{2\mu+1}(g_{p,u})(t)$ preserves sign for $t > 0$ where the function $g_{p,u}$ defines by equality (6.8). Then the function $f(x) := g_{p,u}(x)$, $x \geq 0$, satisfies to the conditions of Theorem 6.5 (for $m = 2\mu + 1$, $f \in H_m^1 \cap H_m$). Moreover, f is continuous at zero and $f(0) = 0$. Therefore, $f(x) = 0$ for almost all $x > 0$. But it is impossible. Therefore (see equality (6.9)) the difference $\frac{1}{\mu((u+\frac{1}{2})^2+t^2)^\mu} - S_\mu(t, u)$ does not preserve sign for $t > 0$ and, consequently, $u^2 + u + \frac{1}{4} < M_\mu(u)$. The statement 1 of Theorem 2.8 proved.

The statement 2 of Theorem 2.8 immediately follows from the equality (6.1).

Let us prove the statement 3. Let $\nu > 0$, $u \geq 0$. Then

$$f'_{\nu,u}(t) = 2t \left((\nu S_\nu(t, u))^{-\frac{1}{\nu}-1} (\nu + 1) S_{\nu+1}(t, u) - 1 \right).$$

The last equality imply that for every $0 < \alpha < \beta$ the next three conditions

$$(6.18a) \quad f'_{\nu,u}(t) \leq (<) 0, \quad \alpha < t < \beta,$$

$$(6.18b) \quad \psi_{\nu,u}(t) := ((\nu + 1) S_{\nu+1}(t, u))^{\frac{1}{\nu+1}} - (\nu S_\nu(t, u))^{\frac{1}{\nu}} \leq (<) 0, \quad \alpha < t < \beta,$$

$$(6.18c) \quad f_{\nu,u}(t) - f_{\nu+1,u}(t) \leq (<) 0, \quad \alpha < t < \beta$$

are equivalent. From the expansion $(1+x)^{\frac{1}{\mu}} = 1 + \frac{x}{\mu} + \frac{x^2(1-\mu)}{2\mu^2} + O(x^3)$, $x \rightarrow 0$, and asymptotic (2.31) follows that the expansion

$$(\mu S_\mu(t, u))^{\frac{1}{\mu}} = \frac{1}{t^2} - \frac{B_2(-u)}{t^4} + \frac{B_4(-u)(\mu+1) + B_2^2(-u)(1-\mu)}{2t^6} + O\left(\frac{1}{t^8}\right), \quad t \rightarrow +\infty$$

takes place for every $\mu > 0$. Therefore,

$$(6.19) \quad \psi_{\nu,u}(t) = \frac{B_4(-u) - B_2^2(-u)}{2t^6} + O\left(\frac{1}{t^8}\right) \sim -\frac{60u^2 + 60u + 11}{360t^6}, \quad t \rightarrow +\infty.$$

This asymptotic implies the existence such a value $t_{\nu,u} \geq 0$ that, for every $t > t_{\nu,u}$, the inequality $\psi_{\nu,u}(t) < 0$ holds and, consequently, the inequality $f'_{\nu,u}(t) < 0$ holds. Therefore, the function $f_{\nu,u}(t)$ strictly decreases with respect to $t \in [t_{\nu,u}, +\infty)$. Consequently, for every $t > t_{\nu,u}$, the inequality $f_{\nu,u}(+\infty) < f_{\nu,u}(t) < f_{\nu,u}(t_{\nu,u})$ holds which is equivalent to the inequality (2.34). The statement **3** of Theorem 2.8 proved.

Let us prove statement **4**. Let, for some $\nu > 0$, $u \geq 0$, the inequality (2.35) be satisfied. From the equivalence of the conditions (6.18), we get that, for every $t > 0$, the inequality $f'_{\nu,u}(t) \leq 0$ be satisfied and, hence, the function $f_{\nu,u}(t)$ decreases on $[0, +\infty)$. Since $f_{\nu,u}(t)$ is analytic function in the neighborhood of every real point then $f_{\nu,u}(t)$ strictly decreases with respect to $t \in [0, +\infty)$ (in the opposite $f'_{\nu,u}(t) = 0$ on some interval and, consequently, for every $t \in \mathbb{R}$ which contradict to the statement **3**). Therefore, the inequality (2.34) be satisfied for $t_{\nu,u} = 0$, $M_\nu(u) = f_{\nu,u}(0)$, and $m_\nu(u) = f_{\nu,u}(+\infty) = u^2 + u + \frac{1}{6}$. From the equivalence of the conditions (6.18), we get, for every $t > 0$, the inequality $f_{\nu,u}(t) \leq f_{\nu+1,u}(t)$ be satisfied and, hence, $m_\nu(u) \leq m_{\nu+1}(u) \leq f_{\nu+1,u}(+\infty) = u^2 + u + \frac{1}{6}$. Therefore, $m_{\nu+1}(u) = u^2 + u + \frac{1}{6}$. If $0 < \mu < \nu + 1$ then the statements **1** and **2** imply the inequality $m_{\nu+1}(u) \leq m_\mu(u) \leq u^2 + u + \frac{1}{6}$ and, hence, $m_\mu(u) = u^2 + u + \frac{1}{6}$.

Let us prove that for $\mu \in (0, \nu+1]$ the inequality $S_\mu(t, u) < \frac{1}{\mu(t^2 + m_\mu(u))^\mu}$ be satisfied for $t \geq 0$. For $\mu = \nu+1$, this inequality follows from the inequality $f_{\nu,u}(t) \leq f_{\nu+1,u}(t)$, $t \geq 0$, which we proved. We used that $m_{\nu+1}(u) = f_{\nu,u}(+\infty) < f_{\nu,u}(t)$, $t \geq 0$. For $0 < \mu < \nu + 1$, we used the equality (6.1) in which $p = m_\mu(u) = u^2 + u + \frac{1}{6}$ and instead ν we put $\nu + 1$.

Let us prove that for every $0 < \mu \leq \nu$, $0 \leq a \leq u^2 + u + \frac{1}{6}$, and $t > 0$, the inequality $\frac{d}{dt}((t^2 + a)^\mu S_\mu(t, u)) > 0$ be satisfied. It is already proved that the inequality (2.34) be satisfied for $t_{\nu,u} = 0$. Therefore, for every $t > 0$, the inequality $\nu S_\nu(t, u) < (t^2 + b)^{-\nu}$, where $b = u^2 + u + \frac{1}{6}$, be satisfied. Taking it and inequality (2.35) into account, we get that for $t > 0$ the inequality

$$(\nu + 1)S_{\nu+1}(t, u) \leq (\nu S_\nu(t, u))^{1+\frac{1}{\nu}} < \frac{\nu S_\nu(t, u)}{t^2 + b}$$

be satisfied. The last inequality is equivalent (see (6.15)) to the inequality

$$\frac{d}{dt}((t^2 + b)^\nu S_\nu(t, u)) > 0, \quad t > 0.$$

Further, we apply the Lemma 6.4. The statement **4** of Theorem 2.8 proved.

The statement **5** immediately follows from the statement **4**, because the inequality (2.35) proved in 1998, for $\nu = 1$, $u = 0$, by Wilkins [13].

The Theorem 2.8 proved. $\square$

Proof of the Theorem 2.9. Let us first prove the statement **1**. It is easy to see that for $p \geq 0$, $\mu > 0$, the next equality

$$(6.20) \quad (-1)^k \psi_{p,u,\mu}^{(k)}(t) = \frac{\Gamma(\mu + k + 1)}{\Gamma(\mu + 1)} \psi_{p,u,\mu+k}(t), \quad k \in \mathbb{Z}_+, \quad t > 0$$

holds. If, for some $\mu > 0$, $\psi_{p,u,\mu} \in M(0, +\infty)$ then $p \geq 0$ and it follows from (6.20) that the inequality $\psi_{p,u,\mu+k}(t) \geq 0$ holds for every $k \in \mathbb{Z}_+$ and $t > 0$. Each of these inequalities is equivalent to the inequality $p \leq m_{\mu+k}(u)$. Passing to the limit in the last inequality as $k \rightarrow +\infty$, we get $p \leq m_\infty(u)$.

If $0 \leq p \leq m_\infty(u)$ then $\psi_{p,u,\mu}(t) \geq 0$ for $\mu > 0$, $t > 0$ and consequently, by the equality (6.20), $\psi_{p,u,\mu} \in M(0, +\infty)$ for $\mu > 0$. The statement **1** proved.

In an analogous way we prove the statement **2**.

Let us prove the statement **3**. It is easy to check that for $p \geq 0$, $u \geq 0$, $\mu > 0$ the next equality

$$\psi_{p,u,\mu}(t) = \frac{1}{\Gamma(\mu + 1)} \int_0^{+\infty} e^{-tx} x^{\mu-1} (e^{-px} - \varphi_u(x)) dx, \quad t > 0$$

holds. From this inequality and the Theorem Hausdorff-Bernstein-Widder follows that, for $p \geq 0$, $u \geq 0$, $\mu > 0$, the condition $\psi_{p,u,\mu} \in M(0, +\infty)$ is equivalent to the inequality $e^{-px} - \varphi_u(x) \geq 0$, $x > 0$. The last inequality is equivalent to $p \leq -\sup_{x>0} \frac{\ln \varphi_u(x)}{x}$. Taking into the statement **1**, the last inequality implies the equality $m_\infty(u) = -\sup_{x>0} \frac{\ln \varphi_u(x)}{x}$. By the same way, we can prove that $M_\infty(u) = -\inf_{x>0} \frac{\ln \varphi_u(x)}{x}$.

Let us prove the statement **4**. The equality (2.28) imply that, for $u \geq 0$ and $x > 0$, the strict inequalities $u^2 + u < -\frac{\ln \varphi_u(x)}{x} < (u+1)^2$ holds. It is obviously that, for every fixed $u \geq 0$, the relation $\varphi_u(x) = 2x(u+1)e^{-x(u+1)^2}(1 + o(1))$, $x \rightarrow +\infty$, holds. This relation and the Example 2.1 imply that next equalities

$$\lim_{x \rightarrow +0} -\frac{\ln \varphi_u(x)}{x} = u^2 + u + \frac{1}{6}, \quad \lim_{x \rightarrow +\infty} -\frac{\ln \varphi_u(x)}{x} = (u+1)^2$$

hold. Therefore,

$$m_\infty(u) = \inf_{x>0} -\frac{\ln \varphi_u(x)}{x} > u^2 + u, \quad m_\infty(u) \leq u^2 + u + \frac{1}{6},$$

$$(u+1)^2 = \lim_{x \rightarrow +\infty} -\frac{\ln \varphi_u(x)}{x} \leq M_\infty(u) = \sup_{x>0} -\frac{\ln \varphi_u(x)}{x} \leq (u+1)^2.$$

The Theorem 2.9 proved. $\square$

Proof of the Theorem 2.11. Let us prove the sufficiency in the statement **1**.

Let $u \geq 0$ and $0 \leq p \leq \sqrt{m_\infty(u)}$. From the Theorem 2.9 follows that, $m_\infty(u) > 0$. The equality (6.9) and right hand side of (2.33) imply that, for every $\mu > 0$ and $0 < p \leq \sqrt{m_\infty(u)}$, the inequality $\mathfrak{F}_{2\mu+1}(g_{p,u})(t) \geq 0$, $t \geq 0$, holds. Next, we use the connection (6.7) between Hankel's transform and Fourier's transform of radial functions. Then, by the Bochner's Theorem, we obtain $g_{p,u} \in \Phi(l_2^{m+1})$, for each $m \in \mathbb{N}$ and $0 < p \leq \sqrt{m_\infty(u)}$. The convergence $g_{p,u} \rightarrow g_{0,u}$, as $p \rightarrow +0$, implies $g_{0,u} \in \Phi(l_2^{m+1})$, for every $m \in \mathbb{N}$. Then, by Schoenberg's Theorem, $g_{p,u} \in \Phi(l_2)$ for every $0 \leq p \leq \sqrt{m_\infty(u)}$. The sufficiency in the statement **1** proved.

Let us prove the necessity in the statement **1**. Let $g_{p,u} \in \Phi(l_2)$. Then, by Schoenberg's Theorem, $g_{p,u} \in \Phi(l_2^{m+1})$, for every $m \in \mathbb{N}$. Boundedness of every positive

definite function implies $p \geq 0$. If $p > 0$ then, by Bochner's Theorem and equality (6.7), the inequality $\mathfrak{F}_{2\mu_m+1}(g_{p,u})(t) \geq 0$, where $t \geq 0$, holds, for every $\mu_m = \frac{m}{2}$, $m \in \mathbb{N}$. The equality (6.9) and right hand side of (2.33) imply $p \leq \sqrt{m_{\mu_m}(u)}$, $m \in \mathbb{N}$. Passing to the limit, as $m \rightarrow \infty$, and taking into account (2.36), we get $p \leq \sqrt{m_\infty(u)}$. The necessity of statement **1** proved.

We can prove the statement **2** in the same way that the statement **1** does. We consider the left hand side inequality in (2.33) instead the right hand side in (2.33).

The statement **3** follows from the next equivalent statements:

- a) The inequality $\frac{d}{dt} \{t^{2\mu} S_\mu(t, u)\} \geq 0$ holds, for every $t > 0$ and $\mu > 0$.
- b) The inequality $\frac{d}{dt} \{t^{2\mu} S_\mu(t, u)\} \geq 0$ holds, for every $t > 0$ and $\mu = \frac{m}{2}$, $m \in \mathbb{N}$.
- c) The inequality $-\mathfrak{F}_{2\mu+1}(h'_u)(t) \geq 0$ holds, for every $t > 0$ and $\mu = \frac{m}{2}$, $m \in \mathbb{N}$.
- d) $-\frac{d}{dx}(h_u(x)) \in \Phi(l_2^{m+1})$, $m \in \mathbb{N}$.
- e) $-\frac{d}{dx}(h_u(x)) \in \Phi(l_2)$.

The Lemma 6.4 implies the equivalence the statements **a** and **b**. The equality (6.11) implies the equivalence of **b** and **c**. The equality (6.7) and Bochner's Theorem implies the equivalence of **c** and **d**. The Schoenberg's Theorem implies the equivalence of **d** and **e**.

The Theorem 2.11 proved. $\square$

REFERENCES

- [1] É.L. Mathieu, *Traité de Physique Mathématique. VI-VII: Théory de l'Élasticité des Corps Solides (Part 2)*, Gauthier-Villars, Paris, 1890.
- [2] L. Berg, "Über eine Abschätzung von Mathieu", *Math. Nachr.*, **7** (1952), 257–259.
- [3] K. Schröder, "Das Problem der eingespannten rechteckigen elastischen Platte", *Math. Ann.*, **121** (1949), 247–326.
- [4] J.G. van der Corput, L.O. Heflinger, "On the inequality of Mathieu", *Indagationes Mathematicae*, **18** (1956), 15–20.
- [5] O. Emersleben, "Über die Reihe $\sum k(k^2 + c^2)^{-2}$ ", *Math. Ann.*, **125** (1952), 165–171.
- [6] G. Polya, "Über die Nullstellen gewisser ganzer Funktionen", *Math. Z.*, **12** (1918), 352–383.
- [7] E. Lukacs, *Characteristic functions*, Griffin, London, 2nd edition, 1970.
- [8] V.P. Zastavnyi, "Extension of a Function from the Exterior of an Interval to a Positive-Definite Function on the Entire Axis and an Approximation Characteristic of the Class $W_M^{r,\beta}$ ", *Ukrainian Mathematical Journal*, **55**(7) (2003), 1189–1197.
- [9] E. Makai, "On the inequality of Mathieu", *Publ. Math. Debrecen*, **5** (1957), 204–205.
- [10] A. Elbert, "Asymptotic expansion and continued fraction for Mathieu's series", *Period. Math. Hungar.*, **13** (1982), 1–8.
- [11] H. Alzer, J.L. Brenner, O.G. Ruehr, "On Mathieu's inequality", *J. Math. Anal. Appl.*, **218** (1998), 607–610.
- [12] H. Alzer, J.L. Brenner, "Problem 97-1", *SIAM Rev.*, **39** (1997), 123.
- [13] J.E. Wilkins, Jr, "An Inequality", *SIAM Rev.*, **40**(1) (1998), 126–128.
- [14] C.L. Wang, X.H. Wang, "A refinement of the Mathieu inequality", *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, No **716-734** (1981) 22–24.
- [15] P.H. Diananda, "Some Inequalities Related to an Inequality of Mathieu", *Math. Ann.*, **250** (1980), 95–98.
- [16] A. Hoorfar, F. Qi, "Some new bounds for Mathieu's series", *Abstract and Applied Analysis*, **2007** (2007), Article ID 94854, 10 pages doi:10.1155/2007/94854.

- [17] F. Qi, CH.-P. Chen, B.-N. Guo, “Notes on double inequalities of Mathieu’s series”, *International Journal of Mathematics and Mathematical Sciences*, **16** (2005), 2547–2554.
- [18] Ž. Tomovski, “New double inequalities for Mathieu type series”, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat.*, **15** (2004), 79–83.
- [19] H. Bateman, A. Erdélyi, *Higher transcendental functions, vol. 1, 2*, New York, Toronto, London, MC Graw-Hill Book Company, INC, 1953.
- [20] T.K. Pogány, H.M. Srivastava, Ž. Tomovski, “Some families of Mathieu **a**-series and alternating Mathieu **a**-series”, *Applied Mathematics and Computation*, **173** (2006), 69–108.
- [21] V.P. Zastavnyi, “On positive definiteness of some functions”, *Journal of Multivariate Analysis*, **73** (2000), 55–81.
- [22] R. M. Trigub, E. S. Belinsky, *Fourier Analysis and Approximation of Functions*, Boston, Dordrecht, London, Kluwer-Springer, 2004.
- [23] N.N. Vakhania, W.I. Tareladze, S.A. Chobanjan, *Probability distributions on Banach Spaces*, Moscow, Nauka, 1985 (in Russian).
- [24] W. Feller, “Completely monotone functions and sequences”, *Trans. Amer. Math. Soc.*, **5** (1939), 662–774.
- [25] I. J. Schoenberg, “Metric spaces and completely monotone functions”, *Ann. Math.*, **39** (1938), 811–841.
- [26] I. J. Schoenberg, “Metric spaces and positive definite functions”, *Trans. Amer. Math. Soc.*, **44** (1938), 522–536.
- [27] M. V. Fedoryuk, *Asymptotics: Integrals and series*, Moscow, Nauka, 1987 (in Russian).
- [28] N.G. de Bruijn, *Asymptotic methods in analysis*, North-Holland Publishing Co., Amsterdam, 1958.
- [29] H. Bateman, A. Erdélyi, W. Magnus, F. Oberhettinger, F. Tricomi, *Tables of integral transforms, vol. I*, New York, Toronto, London, MC Graw-Hill Book Company, INC, 1954.
- [30] V.P. Zastavnyi “Sufficient conditions of an integrability of the Hankel transformation”, *Transactions of IAMM of NAS of Ukraine*, **9** (2004), 68–75 (in Russian).

DEPARTMENT OF MATHEMATICS, DONETSK NATIONAL UNIVERSITY, UNIVERSITETSKAYA STR. 24,
340001, DONETSK, UKRAINE

E-mail address: zastavn@rambler.ru